

C H A P T E R 4

Integration

Section 4.1	Antiderivatives and Indefinite Integration	177
Section 4.2	Area	182
Section 4.3	Riemann Sums and Definite Integrals	188
Section 4.4	The Fundamental Theorem of Calculus	192
Section 4.5	Integration by Substitution	197
Section 4.6	Numerical Integration	204
Review Exercises		209
Problem Solving		214

C H A P T E R 4

Integration

Section 4.1 Antiderivatives and Indefinite Integration

Solutions to Odd-Numbered Exercises

1. $\frac{d}{dx} \left(\frac{3}{x^3} + C \right) = \frac{d}{dx} (3x^{-3} + C) = -9x^{-4} = \frac{-9}{x^4}$

3. $\frac{d}{dx} \left(\frac{1}{3}x^3 - 4x + C \right) = x^2 - 4 = (x - 2)(x + 2)$

5. $\frac{dy}{dt} = 3t^2$

$$y = t^3 + C$$

Check: $\frac{d}{dt}[t^3 + C] = 3t^2$

7. $\frac{dy}{dx} = x^{3/2}$

$$y = \frac{2}{5}x^{5/2} + C$$

Check: $\frac{d}{dx} \left[\frac{2}{5}x^{5/2} + C \right] = x^{3/2}$

<u>Given</u>	<u>Rewrite</u>	<u>Integrate</u>	<u>Simplify</u>
9. $\int \sqrt[3]{x} dx$	$\int x^{1/3} dx$	$\frac{x^{4/3}}{4/3} + C$	$\frac{3}{4}x^{4/3} + C$
11. $\int \frac{1}{x\sqrt{x}} dx$	$\int x^{-3/2} dx$	$\frac{x^{-1/2}}{-1/2} + C$	$-\frac{2}{\sqrt{x}} + C$
13. $\int \frac{1}{2x^3} dx$	$\frac{1}{2} \int x^{-3} dx$	$\frac{1}{2} \left(\frac{x^{-2}}{-2} \right) + C$	$-\frac{1}{4x^2} + C$
15. $\int (x + 3)dx = \frac{x^2}{2} + 3x + C$			17. $\int (2x - 3x^2)dx = x^2 - x^3 + C$
Check: $\frac{d}{dx} \left[\frac{x^2}{2} + 3x + C \right] = x + 3$			Check: $\frac{d}{dx} [x^2 - x^3 + C] = 2x - 3x^2$
19. $\int (x^3 + 2) dx = \frac{1}{4}x^4 + 2x + C$			21. $\int (x^{3/2} + 2x + 1) dx = \frac{2}{5}x^{5/2} + x^2 + x + C$
Check: $\frac{d}{dx} \left(\frac{1}{4}x^4 + 2x + C \right) = x^3 + 2$			Check: $\frac{d}{dx} \left(\frac{2}{5}x^{5/2} + x^2 + x + C \right) = x^{3/2} + 2x + 1$
23. $\int \sqrt[3]{x^2} dx = \int x^{2/3} dx = \frac{x^{5/3}}{5/3} + C = \frac{3}{5}x^{5/3} + C$			25. $\int \frac{1}{x^3} dx = \int x^{-3} dx = \frac{x^{-2}}{-2} + C = -\frac{1}{2x^2} + C$
Check: $\frac{d}{dx} \left(\frac{3}{5}x^{5/3} + C \right) = x^{2/3} = \sqrt[3]{x^2}$			Check: $\frac{d}{dx} \left(-\frac{1}{2x^2} + C \right) = \frac{1}{x^3}$

27. $\int \frac{x^2 + x + 1}{\sqrt{x}} dx = \int (x^{3/2} + x^{1/2} + x^{-1/2}) dx = \frac{2}{5}x^{5/2} + \frac{2}{3}x^{3/2} + 2x^{1/2} + C = \frac{2}{15}x^{1/2}(3x^2 + 5x + 15) + C$

Check: $\frac{d}{dx}\left(\frac{2}{5}x^{5/2} + \frac{2}{3}x^{3/2} + 2x^{1/2} + C\right) = x^{3/2} + x^{1/2} + x^{-1/2} = \frac{x^2 + x + 1}{\sqrt{x}}$

29. $\int (x+1)(3x-2) dx = \int (3x^2 + x - 2) dx$
 $= x^3 + \frac{1}{2}x^2 - 2x + C$

Check: $\frac{d}{dx}\left(x^3 + \frac{1}{2}x^2 - 2x + C\right) = 3x^2 + x - 2$
 $= (x+1)(3x-2)$

31. $\int y^2 \sqrt{y} dy = \int y^{5/2} dy = \frac{2}{7}y^{7/2} + C$

Check: $\frac{d}{dy}\left(\frac{2}{7}y^{7/2} + C\right) = y^{5/2} = y^2 \sqrt{y}$

33. $\int dx = \int 1 dx = x + C$

Check: $\frac{d}{dx}(x + C) = 1$

35. $\int (2 \sin x + 3 \cos x) dx = -2 \cos x + 3 \sin x + C$

Check: $\frac{d}{dx}(-2 \cos x + 3 \sin x + C) = 2 \sin x + 3 \cos x$

37. $\int (1 - \csc t \cot t) dt = t + \csc t + C$

Check: $\frac{d}{dt}(t + \csc t + C) = 1 - \csc t \cot t$


39. $\int (\sec^2 \theta - \sin \theta) d\theta = \tan \theta + \cos \theta + C$

Check: $\frac{d}{d\theta}(\tan \theta + \cos \theta + C) = \sec^2 \theta - \sin \theta$

41. $\int (\tan^2 y + 1) dy = \int \sec^2 y dy = \tan y + C$


Check: $\frac{d}{dy}(\tan y + C) = \sec^2 y = \tan^2 y + 1$

43. $f(x) = \cos x$



45. $f'(x) = 2$


$f(x) = 2x + C$



Answers will vary.

47. $f'(x) = 1 - x^2$

$f(x) = x - \frac{x^3}{3} + C$



Answers will vary.

49. $\frac{dy}{dx} = 2x - 1, (1, 1)$

$y = \int (2x - 1) dx = x^2 - x + C$

$1 = (1)^2 - (1) + C \quad C = 1$
 $y = x^2 - x + 1$


51. $\frac{dy}{dx} = \cos x, (0, 4)$

$$y = \int \cos x \, dx = \sin x + C$$

$$4 = \sin 0 + C \quad C = 4$$

$$y = \sin x + 4$$

53. (a) Answers will vary.




(b) $\frac{dy}{dx} = \frac{1}{2}x - 1, (4, 2)$

$$y = \frac{x^2}{4} - x + C$$

$$2 = \frac{4^2}{4} - 4 + C$$

$$2 = C$$

$$y = \frac{x^2}{4} - x + 2$$



55. $f'(x) = 4x, f(0) = 6$

$$f(x) = \int 4x \, dx = 2x^2 + C$$

$$f(0) = 6 = 2(0)^2 + C \quad C = 6$$

$$f(x) = 2x^2 + 6$$

57. $h'(t) = 8t^3 + 5, h(1) = -4$

$$h(t) = \int (8t^3 + 5) dt = 2t^4 + 5t + C$$

$$h(1) = -4 = 2 + 5 + C \quad C = -11$$

$$h(t) = 2t^4 + 5t - 11$$

59. $f''(x) = 2$

$$f'(2) = 5$$

$$f(2) = 10$$

$$f'(x) = \int 2 \, dx = 2x + C_1$$

$$f'(2) = 4 + C_1 = 5 \quad C_1 = 1$$

$$f'(x) = 2x + 1$$

$$f(x) = \int (2x + 1) \, dx = x^2 + x + C_2$$

$$f(2) = 6 + C_2 = 10 \quad C_2 = 4$$

$$f(x) = x^2 + x + 4$$

61. $f''(x) = x^{-3/2}$

$$f'(4) = 2$$

$$f(0) = 0$$

$$f'(x) = \int x^{-3/2} \, dx = -2x^{-1/2} + C_1 = -\frac{2}{\sqrt{x}} + C_1$$

$$f'(4) = -\frac{2}{2} + C_1 = 2 \quad C_1 = 3$$

$$f'(x) = -\frac{2}{\sqrt{x}} + 3$$

$$f(x) = \int (-2x^{-1/2} + 3) \, dx = -4x^{1/2} + 3x + C_2$$

$$f(0) = 0 + 0 + C_2 = 0 \quad C_2 = 0$$

$$f(x) = -4x^{1/2} + 3x = -4\sqrt{x} + 3x$$

63. (a) $h(t) = \int (1.5t + 5) \, dt = 0.75t^2 + 5t + C$

$$h(0) = 0 + 0 + C = 12 \quad C = 12$$

$$h(t) = 0.75t^2 + 5t + 12$$

(b) $h(6) = 0.75(6)^2 + 5(6) + 12 = 69 \text{ cm}$

65. $f(0) = -4$. Graph of f' is given.

- (a) $f'(4) \approx -1.0$
- (b) No. The slopes of the tangent lines are greater than 2 on $[0, 2]$. Therefore, f must increase more than 4 units on $[0, 4]$.
- (c) No, $f(5) < f(4)$ because f is decreasing on $[4, 5]$.
- (d) f has a maximum at $x = 3.5$ because $f'(3.5) \approx 0$ and the first derivative test.
- (e) f is concave upward when f' is increasing on $(-\infty, 1)$ and $(5, \infty)$. f is concave downward on $(1, 5)$. Points of inflection at $x = 1, 5$.

67. $a(t) = -32 \text{ ft/sec}^2$

$$v(t) = \int -32 dt = -32t + C_1$$

$$v(0) = 60 = C_1$$

$$s(t) = \int (-32t + 60)dt = -16t^2 + 60t + C_2$$

$$s(0) = 6 = C_2$$

$s(t) = -16t^2 + 60t + 6$ Position function

The ball reaches its maximum height when

$$v(t) = -32t + 60 = 0$$

$$32t = 60$$

$$t = \frac{15}{8} \text{ seconds}$$

$$s\left(\frac{15}{8}\right) = -16\left(\frac{15}{8}\right)^2 + 60\left(\frac{15}{8}\right) + 6 \approx 62.26 \text{ feet}$$

71. $a(t) = -9.8$

$$v(t) = \int -9.8 dt = -9.8t + C_1$$


$$v(0) = v_0 = C_1 \quad v(t) = -9.8t + v_0$$

$$f(t) = \int (-9.8t + v_0) dt = -4.9t^2 + v_0 t + C_2$$

$$f(0) = s_0 = C_2 \quad f(t) = -4.9t^2 + v_0 t + s_0$$

- (f) f'' is a minimum at $x = 3$.

(g)



69. From Exercise 68, we have:

$$s(t) = -16t^2 + v_0 t$$

$$s'(t) = -32t + v_0 = 0 \text{ when } t = \frac{v_0}{32} = \text{time to reach maximum height.}$$

$$s\left(\frac{v_0}{32}\right) = -16\left(\frac{v_0}{32}\right)^2 + v_0\left(\frac{v_0}{32}\right) = 550$$

$$-\frac{v_0^2}{64} + \frac{v_0^2}{32} = 550$$

$$v_0^2 = 35,200$$

$$v_0 \approx 187.617 \text{ ft/sec}$$

73. From Exercise 71, $f(t) = -4.9t^2 + 10t + 2$.

$$v(t) = -9.8t + 10 = 0 \text{ (Maximum height when } v = 0.)$$

$$9.8t = 10$$

$$t = \frac{10}{9.8}$$

$$f\left(\frac{10}{9.8}\right) \approx 7.1 \text{ m}$$

75. $a = -1.6$

$$v(t) = \int -1.6 dt = -1.6t + v_0 = -1.6t, \text{ since the stone was dropped, } v_0 = 0.$$

$$s(t) = \int (-1.6t) dt = -0.8t^2 + s_0$$

$$s(20) = 0 \quad -0.8(20)^2 + s_0 = 0$$

$$s_0 = 320$$

Thus, the height of the cliff is 320 meters.

$$v(t) = -1.6t$$

$$v(20) = -32 \text{ m/sec}$$

77. $x(t) = t^3 - 6t^2 + 9t - 2 \quad 0 \leq t \leq 5$

$$\begin{aligned} \text{(a)} \quad v(t) &= x'(t) = 3t^2 - 12t + 9 \\ &= 3(t^2 - 4t + 3) = 3(t-1)(t-3) \\ a(t) &= v'(t) = 6t - 12 = 6(t-2) \\ \text{(b)} \quad v(t) &> 0 \text{ when } 0 < t < 1 \text{ or } 3 < t < 5. \\ \text{(c)} \quad a(t) &= 6(t-2) = 0 \text{ when } t = 2. \\ v(2) &= 3(1)(-1) = -3 \end{aligned}$$

81. (a) $v(0) = 25 \text{ km/hr} = 25 \cdot \frac{1000}{3600} = \frac{250}{36} \text{ m/sec}$

$$v(13) = 80 \text{ km/hr} = 80 \cdot \frac{1000}{3600} = \frac{800}{36} \text{ m/sec}$$

$a(t) = a$ (constant acceleration)

$$v(t) = at + C$$

$$v(0) = \frac{250}{36} \quad v(t) = at + \frac{250}{36}$$

$$v(13) = \frac{800}{36} = 13a + \frac{250}{36}$$

$$\frac{550}{36} = 13a$$

$$a = \frac{550}{468} = \frac{275}{234} \approx 1.175 \text{ m/sec}^2$$

(b) $s(t) = a\frac{t^2}{2} + \frac{250}{36}t \quad (s(0) = 0)$

$$s(13) = \frac{275}{234} \frac{(13)^2}{2} + \frac{250}{36}(13) \approx 189.58 \text{ m}$$

85. $\frac{(1 \text{ mi/hr})(5280 \text{ ft/mi})}{(3600 \text{ sec/hr})} = \frac{22}{15} \text{ ft/sec}$

(a)	<table border="1"> <thead> <tr> <th>t</th><th>0</th><th>5</th><th>10</th><th>15</th><th>20</th><th>25</th><th>30</th></tr> </thead> <tbody> <tr> <td>$V_1(\text{ft/sec})$</td><td>0</td><td>3.67</td><td>10.27</td><td>23.47</td><td>42.53</td><td>66</td><td>95.33</td></tr> <tr> <td>$V_2(\text{ft/sec})$</td><td>0</td><td>30.8</td><td>55.73</td><td>74.8</td><td>88</td><td>93.87</td><td>95.33</td></tr> </tbody> </table>	t	0	5	10	15	20	25	30	$V_1(\text{ft/sec})$	0	3.67	10.27	23.47	42.53	66	95.33	$V_2(\text{ft/sec})$	0	30.8	55.73	74.8	88	93.87	95.33
t	0	5	10	15	20	25	30																		
$V_1(\text{ft/sec})$	0	3.67	10.27	23.47	42.53	66	95.33																		
$V_2(\text{ft/sec})$	0	30.8	55.73	74.8	88	93.87	95.33																		

(c) $S_1(t) = \int V_1(t) dt = \frac{0.1068}{3}t^3 - \frac{0.0416}{2}t^2 + 0.3679t$

$$S_2(t) = \int V_2(t) dt = -\frac{0.1208}{3}t^3 + \frac{6.7991}{2}t^2 - 0.0707t$$

[In both cases, the constant of integration is 0 because $S_1(0) = S_2(0) = 0$]

$$S_1(30) \approx 953.5 \text{ feet}$$

$$S_2(30) \approx 1970.3 \text{ feet}$$

The second car was going faster than the first until the end.

79. $v(t) = \frac{1}{\sqrt{t}} = t^{-1/2} \quad t > 0$

$$x(t) = \int v(t) dt = 2t^{1/2} + C$$

$$x(1) = 4 = 2(1) + C \quad C = 2$$

$x(t) = 2t^{1/2} + 2$ position function

$$a(t) = v'(t) = -\frac{1}{2}t^{-3/2} = \frac{-1}{2t^{3/2}} \text{ acceleration}$$

83. Truck: $v(t) = 30$

$$s(t) = 30t \text{ (Let } s(0) = 0.)$$

Automobile: $a(t) = 6$

$$v(t) = 6t \text{ (Let } v(0) = 0.)$$

$$s(t) = 3t^2 \text{ (Let } s(0) = 0.)$$

At the point where the automobile overtakes the truck:

$$30t = 3t^2$$

$$0 = 3t^2 - 30t$$

$$0 = 3t(t-10) \text{ when } t = 10 \text{ sec.}$$

(a) $s(10) = 3(10)^2 = 300 \text{ ft}$

(b) $v(10) = 6(10) = 60 \text{ ft/sec} \approx 41 \text{ mph}$

(b) $V_1(t) = 0.1068t^2 - 0.0416t + 0.3679$

$$V_2(t) = -0.1208t^3 + 6.7991t - 0.0707$$

87. $a(t) = k$

$$v(t) = kt$$

$$s(t) = \frac{k}{2}t^2 \text{ since } v(0) = s(0) = 0.$$

At the time of lift-off, $kt = 160$ and $(k/2)t^2 = 0.7$. Since $(k/2)t^2 = 0.7$,

$$t = \sqrt{\frac{1.4}{k}}$$

$$v\left(\sqrt{\frac{1.4}{k}}\right) = k\sqrt{\frac{1.4}{k}} = 160$$

$$1.4k = 160^2 \quad k = \frac{160^2}{1.4}$$

$$\approx 18,285.714 \text{ mi/hr}^2$$

$$\approx 7.45 \text{ ft/sec}^2.$$

89. True

91. True

93. False. For example, $\int x \cdot x dx \neq \int x dx \cdot \int x dx$ because $\frac{x^3}{3} + C \neq \left(\frac{x^2}{2} + C_1\right)\left(\frac{x^2}{2} + C_2\right)$

95. $f'(x) = \begin{cases} 1, & 0 < x < 2 \\ 3x, & 2 < x < 5 \end{cases}$

$$f(x) = \begin{cases} x + C_1, & 0 < x < 2 \\ \frac{3x^2}{2} + C_2, & 2 < x < 5 \end{cases}$$

$$f(1) = 3 \quad 1 + C_1 = 3 \quad C_1 = 2$$

f is continuous: Values must agree at $x = 2$:

$$4 = 6 + C_2 \quad C_2 = -2$$

$$f(x) = \begin{cases} x + 2, & 0 < x < 2 \\ \frac{3x^2}{2} - 2, & 2 < x < 5 \end{cases}$$

The left and right hand derivatives at $x = 2$ do not agree. Hence f is not differentiable at $x = 2$.

Section 4.2 Area

1. $\sum_{i=1}^5 (2i + 1) = 2\sum_{i=1}^5 i + \sum_{i=1}^5 1 = 2(1 + 2 + 3 + 4 + 5) + 5 = 35$

3. $\sum_{k=0}^4 \frac{1}{k^2 + 1} = 1 + \frac{1}{2} + \frac{1}{5} + \frac{1}{10} + \frac{1}{17} = \frac{158}{85}$

5. $\sum_{k=1}^4 c = c + c + c + c = 4c$

7. $\sum_{i=1}^9 \frac{1}{3i}$

9. $\sum_{j=1}^8 \left[5\left(\frac{j}{8}\right) + 3 \right]$

11. $\frac{2}{n} \sum_{i=1}^n \left[\left(\frac{2i}{n}\right)^3 - \left(\frac{2i}{n}\right) \right]$

13. $\frac{3}{n} \sum_{i=1}^n \left[2\left(1 + \frac{3i}{n}\right)^2 \right]$

15. $\sum_{i=1}^{20} 2i = 2\sum_{i=1}^{20} i = 2\left[\frac{20(21)}{2}\right] = 420$

17. $\sum_{i=1}^{20} (i - 1)^2 = \sum_{i=1}^{19} i^2$

$$= \left[\frac{19(20)(39)}{6} \right] = 2470$$

$$\begin{aligned}
19. \sum_{i=1}^{15} i(i-1)^2 &= \sum_{i=1}^{15} i^3 - 2 \sum_{i=1}^{15} i^2 + \sum_{i=1}^{15} i \\
&= \frac{15^2(16)^2}{4} - 2 \frac{15(16)(31)}{6} + \frac{15(16)}{2} \\
&= 14,400 - 2,480 + 120 \\
&= 12,040
\end{aligned}$$

$$\begin{aligned}
23. S &= [3 + 4 + \frac{9}{2} + 5](1) = \frac{33}{2} = 16.5 \\
s &= [1 + 3 + 4 + \frac{9}{2}](1) = \frac{25}{2} = 12.5
\end{aligned}$$

$$\begin{aligned}
27. S(4) &= \sqrt{\frac{1}{4}}\left(\frac{1}{4}\right) + \sqrt{\frac{1}{2}}\left(\frac{1}{4}\right) + \sqrt{\frac{3}{4}}\left(\frac{1}{4}\right) + \sqrt{1}\left(\frac{1}{4}\right) = \frac{1 + \sqrt{2} + \sqrt{3} + 2}{8} \approx 0.768 \\
s(4) &= 0\left(\frac{1}{4}\right) + \sqrt{\frac{1}{4}}\left(\frac{1}{4}\right) + \sqrt{\frac{1}{2}}\left(\frac{1}{4}\right) + \sqrt{\frac{3}{4}}\left(\frac{1}{4}\right) = \frac{1 + \sqrt{2} + \sqrt{3}}{8} \approx 0.518
\end{aligned}$$

$$\begin{aligned}
29. S(5) &= 1\left(\frac{1}{5}\right) + \frac{1}{6/5}\left(\frac{1}{5}\right) + \frac{1}{7/5}\left(\frac{1}{5}\right) + \frac{1}{8/5}\left(\frac{1}{5}\right) + \frac{1}{9/5}\left(\frac{1}{5}\right) = \frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8} + \frac{1}{9} \approx 0.746 \\
s(5) &= \frac{1}{6/5}\left(\frac{1}{5}\right) + \frac{1}{7/5}\left(\frac{1}{5}\right) + \frac{1}{8/5}\left(\frac{1}{5}\right) + \frac{1}{9/5}\left(\frac{1}{5}\right) + \frac{1}{2}\left(\frac{1}{5}\right) = \frac{1}{6} + \frac{1}{7} + \frac{1}{8} + \frac{1}{9} + \frac{1}{10} \approx 0.646
\end{aligned}$$

$$31. \lim_{n \rightarrow \infty} \left[\left(\frac{81}{n^4} \right) \frac{n^2(n+1)^2}{4} \right] = \frac{81}{4} \lim_{n \rightarrow \infty} \left[\frac{n^4 + 2n^3 + n^2}{n^4} \right] = \frac{81}{4}(1) = \frac{81}{4}$$

$$33. \lim_{n \rightarrow \infty} \left[\left(\frac{18}{n^2} \right) \frac{n(n+1)}{2} \right] = \frac{18}{2} \lim_{n \rightarrow \infty} \left[\frac{n^2 + n}{n^2} \right] = \frac{18}{2}(1) = 9$$

$$\begin{aligned}
35. \sum_{i=1}^n \frac{2i+1}{n^2} &= \frac{1}{n^2} \sum_{i=1}^n (2i+1) = \frac{1}{n^2} \left[2 \frac{n(n+1)}{2} + n \right] = \frac{n+2}{n} = S(n) \\
S(10) &= \frac{12}{10} = 1.2
\end{aligned}$$

$$S(100) = 1.02$$

$$S(1000) = 1.002$$

$$S(10,000) = 1.0002$$

$$\begin{aligned}
37. \sum_{k=1}^n \frac{6k(k-1)}{n^3} &= \frac{6}{n^3} \sum_{k=1}^n (k^2 - k) = \frac{6}{n^3} \left[\frac{n(n+1)(2n+1)}{6} - \frac{n(n+1)}{2} \right] \\
&= \frac{6}{n^2} \left[\frac{2n^2 + 3n + 1 - 3n - 3}{6} \right] = \frac{1}{n^2}[2n^2 - 2] = S(n)
\end{aligned}$$

$$S(10) = 1.98$$

$$S(100) = 1.9998$$


$$S(1000) = 1.999998$$

$$S(10,000) = 1.99999998$$

$$39. \lim_{n \rightarrow \infty} \sum_{i=1}^n \left(\frac{16i}{n^2} \right) = \lim_{n \rightarrow \infty} \frac{16}{n^2} \sum_{i=1}^n i = \lim_{n \rightarrow \infty} \frac{16}{n^2} \left(\frac{n(n+1)}{2} \right) = \lim_{n \rightarrow \infty} \left[8 \left(\frac{n^2+n}{n^2} \right) \right] = 8 \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n} \right) = 8$$

41. $\lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{1}{n^3}(i-1)^2 = \lim_{n \rightarrow \infty} \frac{1}{n^3} \sum_{i=1}^{n-1} i^2 = \lim_{n \rightarrow \infty} \frac{1}{n^3} \left[\frac{(n-1)(n)(2n-1)}{6} \right]$
 $= \lim_{n \rightarrow \infty} \frac{1}{6} \left[\frac{2n^3 - 3n^2 + n}{n^3} \right] = \lim_{n \rightarrow \infty} \left[\frac{1}{6} \left(\frac{2 - (3/n) + (1/n^2)}{1} \right) \right] = \frac{1}{3}$

43. $\lim_{n \rightarrow \infty} \sum_{i=1}^n \left(1 + \frac{i}{n} \right) \left(\frac{2}{n} \right) = 2 \lim_{n \rightarrow \infty} \frac{1}{n} \left[\sum_{i=1}^n 1 + \frac{1}{n} \sum_{i=1}^n i \right] = 2 \lim_{n \rightarrow \infty} \frac{1}{n} \left[n + \frac{1}{n} \left(\frac{n(n+1)}{2} \right) \right] = 2 \lim_{n \rightarrow \infty} \left[1 + \frac{n^2+n}{2n^2} \right] = 2 \left(1 + \frac{1}{2} \right) = 3$



(b) $\Delta x = \frac{2-0}{n} = \frac{2}{n}$

Endpoints:

$$0 < 1\left(\frac{2}{n}\right) < 2\left(\frac{2}{n}\right) < \dots < (n-1)\left(\frac{2}{n}\right) < n\left(\frac{2}{n}\right) = 2$$

(c) Since $y = x$ is increasing, $f(m_i) = f(x_{i-1})$ on $[x_{i-1}, x_i]$.

$$\begin{aligned} s(n) &= \sum_{i=1}^n f(x_{i-1}) \Delta x \\ &= \sum_{i=1}^n f\left(\frac{2i-2}{n}\right) \left(\frac{2}{n}\right) = \sum_{i=1}^n \left[(i-1)\left(\frac{2}{n}\right) \right] \left(\frac{2}{n}\right) \end{aligned}$$


(d) $f(M_i) = f(x_i)$ on $[x_{i-1}, x_i]$

$$S(n) = \sum_{i=1}^n f(x_i) \Delta x = \sum_{i=1}^n f\left(\frac{2i}{n}\right) \frac{2}{n} = \sum_{i=1}^n \left[i\left(\frac{2}{n}\right) \right] \left(\frac{2}{n}\right)$$

47. $y = -2x + 3$ on $[0, 1]$. $\left(\text{Note: } \Delta x = \frac{1-0}{n} = \frac{1}{n} \right)$

$$\begin{aligned} s(n) &= \sum_{i=1}^n f\left(\frac{i}{n}\right) \left(\frac{1}{n}\right) = \sum_{i=1}^n \left[-2\left(\frac{i}{n}\right) + 3 \right] \left(\frac{1}{n}\right) \\ &= 3 - \frac{2}{n^2} \sum_{i=1}^n i = 3 - \frac{2(n+1)n}{2n^2} = 2 - \frac{1}{n} \end{aligned}$$


$$\text{Area} = \lim_{n \rightarrow \infty} s(n) = 2$$



49. $y = x^2 + 2$ on $[0, 1]$. $\left(\text{Note: } \Delta x = \frac{1}{n} \right)$

$$\begin{aligned} S(n) &= \sum_{i=1}^n f\left(\frac{i}{n}\right) \left(\frac{1}{n}\right) = \sum_{i=1}^n \left[\left(\frac{i}{n}\right)^2 + 2 \right] \left(\frac{1}{n}\right) \\ &= \left[\frac{1}{n^3} \sum_{i=1}^n i^2 \right] + 2 = \frac{n(n+1)(2n+1)}{6n^3} + 2 = \frac{1}{6} \left(2 + \frac{3}{n} + \frac{1}{n^2} \right) + 2 \end{aligned}$$

$$\text{Area} = \lim_{n \rightarrow \infty} S(n) = \frac{7}{3}$$



(e)


x	5	10	50	100
$s(n)$	1.6	1.8	1.96	1.98
$S(n)$	2.4	2.2	2.04	2.02

$$\begin{aligned} (f) \quad &\lim_{n \rightarrow \infty} \sum_{i=1}^n \left[(i-1)\left(\frac{2}{n}\right) \right] \left(\frac{2}{n}\right) = \lim_{n \rightarrow \infty} \frac{4}{n^2} \sum_{i=1}^n (i-1) \\ &= \lim_{n \rightarrow \infty} \frac{4}{n^2} \left[\frac{n(n+1)}{2} - n \right] = \lim_{n \rightarrow \infty} \left[\frac{2(n+1)}{n} - \frac{4}{n} \right] = 2 \\ &\lim_{n \rightarrow \infty} \sum_{i=1}^n \left[i\left(\frac{2}{n}\right) \right] \left(\frac{2}{n}\right) = \lim_{n \rightarrow \infty} \frac{4}{n^2} \sum_{i=1}^n i \\ &= \lim_{n \rightarrow \infty} \left(\frac{4}{n^2} \right) \frac{n(n+1)}{2} = \lim_{n \rightarrow \infty} \frac{2(n+1)}{n} = 2 \end{aligned}$$

51. $y = 16 - x^2$ on $[1, 3]$. (Note: $\Delta x = \frac{2}{n}$)

$$\begin{aligned} s(n) &= \sum_{i=1}^n f\left(1 + \frac{2i}{n}\right)\left(\frac{2}{n}\right) = \sum_{i=1}^n \left[16 - \left(1 + \frac{2i}{n}\right)^2\right]\left(\frac{2}{n}\right) \\ &= \frac{2}{n} \sum_{i=1}^n \left[15 - \frac{4i^2}{n^2} - \frac{4i}{n}\right] \\ &= \frac{2}{n} \left[15n - \frac{4}{n^2} \frac{n(n+1)(2n+1)}{6} - \frac{4}{n} \frac{n(n+1)}{2}\right] \\ &= 30 - \frac{8}{6n^2}(n+1)(2n+1) - \frac{4}{n}(n+1) \end{aligned}$$


$$\text{Area} = \lim_{n \rightarrow \infty} s(n) = 30 - \frac{8}{3} - 4 = \frac{70}{3} = 23\frac{1}{3}$$



53. $y = 64 - x^3$ on $[1, 4]$. (Note: $\Delta x = \frac{3}{n}$)

$$\begin{aligned} s(n) &= \sum_{i=1}^n f\left(1 + \frac{3i}{n}\right)\left(\frac{3}{n}\right) = \sum_{i=1}^n \left[64 - \left(1 + \frac{3i}{n}\right)^3\right]\left(\frac{3}{n}\right) \\ &= \frac{3}{n} \sum_{i=1}^n \left[63 - \frac{27i^3}{n^3} - \frac{27i^2}{n^2} - \frac{9i}{n}\right] \\ &= \frac{3}{n} \left[63n - \frac{27}{n^3} \frac{n^2(n+1)^2}{4} - \frac{27}{n^2} \frac{n(n+1)(2n+1)}{6} - \frac{9}{n} \frac{n(n+1)}{2}\right] \\ &= 189 - \frac{81}{4n^2}(n+1)^2 - \frac{81}{6n^2}(n+1)(2n+1) - \frac{27}{2} \frac{n+1}{n} \end{aligned}$$

$$\text{Area} = \lim_{n \rightarrow \infty} s(n) = 189 - \frac{81}{4} - 27 - \frac{27}{2} = \frac{513}{4} = 128.25$$




55. $y = x^2 - x^3$ on $[-1, 1]$. (Note: $\Delta x = \frac{1 - (-1)}{n} = \frac{2}{n}$)

Again, $T(n)$ is neither an upper nor a lower sum.

$$\begin{aligned} T(n) &= \sum_{i=1}^n f\left(-1 + \frac{2i}{n}\right)\left(\frac{2}{n}\right) = \sum_{i=1}^n \left[\left(-1 + \frac{2i}{n}\right)^2 - \left(-1 + \frac{2i}{n}\right)^3\right]\left(\frac{2}{n}\right) \\ &= \sum_{i=1}^n \left[\left(1 - \frac{4i}{n} + \frac{4i^2}{n^2}\right) - \left(-1 + \frac{6i}{n} - \frac{12i^2}{n^2} + \frac{8i^3}{n^3}\right)\right]\left(\frac{2}{n}\right) \\ &= \sum_{i=1}^n \left[2 - \frac{10i}{n} + \frac{16i^2}{n^2} - \frac{8i^3}{n^3}\right]\left(\frac{2}{n}\right) = \frac{4}{n^2} \sum_{i=1}^n 1 - \frac{20}{n^3} \sum_{i=1}^n i + \frac{32}{n^4} \sum_{i=1}^n i^2 - \frac{16}{n^5} \sum_{i=1}^n i^3 \\ &= \frac{4}{n}(n) - \frac{20}{n^2} \cdot \frac{n(n+1)}{2} + \frac{32}{n^3} \cdot \frac{n(n+1)(2n+1)}{6} - \frac{16}{n^4} \cdot \frac{n^2(n+1)^2}{4} \\ &= 4 - 10\left(1 + \frac{1}{n}\right) + \frac{16}{3}\left(2 + \frac{3}{n} + \frac{1}{n^2}\right) - 4\left(1 + \frac{2}{n} + \frac{1}{n^2}\right) \end{aligned}$$


$$\text{Area} = \lim_{n \rightarrow \infty} T(n) = 4 - 10 + \frac{32}{3} - 4 = \frac{2}{3}$$



57. $f(y) = 3y, 0 \leq y \leq 2$ (**Note:** $\Delta y = \frac{2-0}{n} = \frac{2}{n}$)

$$\begin{aligned} S(n) &= \sum_{i=1}^n f(m_i) \Delta y = \sum_{i=1}^n f\left(\frac{2i}{n}\right)\left(\frac{2}{n}\right) = \sum_{i=1}^n 3\left(\frac{2i}{n}\right)\left(\frac{2}{n}\right) \\ &= \frac{12}{n^2} \sum_{i=1}^n i = \left(\frac{12}{n^2}\right) \cdot \frac{n(n+1)}{2} = \frac{6(n+1)}{n} = 6 + \frac{6}{n} \end{aligned}$$


$$\text{Area} = \lim_{n \rightarrow \infty} S(n) = \lim_{n \rightarrow \infty} \left(6 + \frac{6}{n}\right) = 6$$



59. $f(y) = y^2, 0 \leq y \leq 3$ (**Note:** $\Delta y = \frac{3-0}{n} = \frac{3}{n}$)

$$\begin{aligned} S(n) &= \sum_{i=1}^n f\left(\frac{3i}{n}\right)\left(\frac{3}{n}\right) = \sum_{i=1}^n \left(\frac{3i}{n}\right)^2\left(\frac{3}{n}\right) = \frac{27}{n^3} \sum_{i=1}^n i^2 \\ &= \frac{27}{n^3} \cdot \frac{n(n+1)(2n+1)}{6} = \frac{9}{n^2} \left(\frac{2n^2+3n+1}{2}\right) = 9 + \frac{27}{2n} + \frac{9}{2n^2} \end{aligned}$$


$$\text{Area} = \lim_{n \rightarrow \infty} S(n) = \lim_{n \rightarrow \infty} \left(9 + \frac{27}{2n} + \frac{9}{2n^2}\right) = 9$$



61. $g(y) = 4y^2 - y^3, 1 \leq y \leq 3$. (**Note:** $\Delta y = \frac{3-1}{n} = \frac{2}{n}$)

$$\begin{aligned} S(n) &= \sum_{i=1}^n g\left(1 + \frac{2i}{n}\right)\left(\frac{2}{n}\right) \\ &= \sum_{i=1}^n \left[4\left(1 + \frac{2i}{n}\right)^2 - \left(1 + \frac{2i}{n}\right)^3\right] \frac{2}{n} \\ &= \frac{2}{n} \sum_{i=1}^n 4 \left[1 + \frac{4i}{n} + \frac{4i^2}{n^2}\right] - \left[1 + \frac{6i}{n} + \frac{12i^2}{n^2} + \frac{8i^3}{n^3}\right] \\ &= \frac{2}{n} \sum_{i=1}^n \left[3 + \frac{10i}{n} + \frac{4i^2}{n^2} - \frac{8i^3}{n^3}\right] = \frac{2}{n} \left[3n + \frac{10}{n} \frac{n(n+1)}{2} + \frac{4}{n^2} \frac{n(n+1)(2n+1)}{6} - \frac{8}{n^3} \frac{n^2(n+1)^2}{4}\right] \end{aligned}$$

$$\text{Area} = \lim_{n \rightarrow \infty} S(n) = 6 + 10 + \frac{8}{3} - 4 = \frac{44}{3}$$



63. $f(x) = x^2 + 3, 0 \leq x \leq 2, n = 4$

$$\text{Let } c_i = \frac{x_i + x_{i-1}}{2}.$$

$$\Delta x = \frac{1}{2}, c_1 = \frac{1}{4}, c_2 = \frac{3}{4}, c_3 = \frac{5}{4}, c_4 = \frac{7}{4}$$

$$\begin{aligned} \text{Area} &\approx \sum_{i=1}^n f(c_i) \Delta x = \sum_{i=1}^4 [c_i^2 + 3]\left(\frac{1}{2}\right) \\ &= \frac{1}{2} \left[\left(\frac{1}{16} + 3\right) + \left(\frac{9}{16} + 3\right) + \left(\frac{25}{16} + 3\right) + \left(\frac{49}{16} + 3\right) \right] \\ &= \frac{69}{8} \end{aligned}$$

65. $f(x) = \tan x, 0 \leq x \leq \frac{\pi}{4}, n = 4$

$$\text{Let } c_i = \frac{x_i + x_{i-1}}{2}.$$

$$\Delta x = \frac{\pi}{16}, c_1 = \frac{\pi}{32}, c_2 = \frac{3\pi}{32}, c_3 = \frac{5\pi}{32}, c_4 = \frac{7\pi}{32}$$

$$\begin{aligned} \text{Area} &\approx \sum_{i=1}^n f(c_i) \Delta x = \sum_{i=1}^4 (\tan c_i) \left(\frac{\pi}{16}\right) \\ &= \frac{\pi}{16} \left(\tan \frac{\pi}{32} + \tan \frac{3\pi}{32} + \tan \frac{5\pi}{32} + \tan \frac{7\pi}{32} \right) \approx 0.345 \end{aligned}$$

67. $f(x) = \sqrt{x}$ on $[0, 4]$.


n	4	8	12	16	20
Approximate area	5.3838	5.3523	5.3439	5.3403	5.3384

(Exact value is $16/3$)


69. $f(x) = \tan\left(\frac{\pi x}{8}\right)$ on $[1, 3]$.

n	4	8	12	16	20
Approximate area	2.2223	2.2387	2.2418	2.2430	2.2435

71. We can use the line $y = x$ bounded by $x = a$ and $x = b$.
The sum of the areas of these inscribed rectangles is the lower sum.




The sum of the areas of these circumscribed rectangles is the upper sum.



We can see that the rectangles do not contain all of the area in the first graph and the rectangles in the second graph cover more than the area of the region.

The exact value of the area lies between these two sums.


73. (a)



Lower sum:

$$s(4) = 0 + 4 + 5\frac{1}{3} + 6 = 15\frac{1}{3} = \frac{46}{3} \approx 15.333$$

(c)




Midpoint Rule:

$$M(4) = 2\frac{2}{3} + 4\frac{4}{5} + 5\frac{5}{7} + 6\frac{2}{9} = \frac{6112}{315} \approx 19.403$$

(e)

n	4	8	20	100	200
$s(n)$	15.333	17.368	18.459	18.995	19.06
$S(n)$	21.733	20.568	19.739	19.251	19.188
$M(n)$	19.403	19.201	19.137	19.125	19.125

(b)




Upper sum:

$$S(4) = 4 + 5\frac{1}{3} + 6 + 6\frac{2}{5} = 21\frac{11}{15} = \frac{326}{15} \approx 21.733$$

- (d) In each case, $\Delta x = 4/n$. The lower sum uses left endpoints, $(i-1)(4/n)$. The upper sum uses right endpoints, $(i)(4/n)$. The Midpoint Rule uses midpoints, $(i - \frac{1}{2})(4/n)$.

- (f) $s(n)$ increases because the lower sum approaches the exact value as n increases. $S(n)$ decreases because the upper sum approaches the exact value as n increases. Because of the shape of the graph, the lower sum is always smaller than the exact value, whereas the upper sum is always larger.

75.



77. True. (Theorem 4.2 (2))

b. $A \approx 6$ square units


79. $f(x) = \sin x, \left[0, \frac{\pi}{2}\right]$

Let A_1 = area bounded by $f(x) = \sin x$, the x -axis, $x = 0$ and $x = \pi/2$. Let A_2 = area of the rectangle bounded by $y = 1$, $y = 0$, $x = 0$, and $x = \pi/2$. Thus, $A_2 = (\pi/2)(1) \approx 1.570796$.

In this program, the computer is generating N_2 pairs of random points in the rectangle whose area is represented by A_2 . It is keeping track of how many of these points, N_1 , lie in the region whose area is represented by A_1 . Since the points are randomly generated, we assume that

$$\frac{A_1}{A_2} \approx \frac{N_1}{N_2} \quad A_1 \approx \frac{N_1}{N_2} A_2.$$

The larger N_2 is the better the approximation to A_1 .



81. Suppose there are n rows in the figure. The stars on the left total $1 + 2 + \dots + n$, as do the stars on the right. There are $n(n + 1)$ stars in total, hence


$$2[1 + 2 + \dots + n] = n(n + 1)$$

$$1 + 2 + \dots + n = \frac{1}{2}(n)(n + 1).$$

83. (a) $y = (-4.09 \times 10^{-5})x^3 + 0.016x^2 - 2.67x + 452.9$

- (c) Using the integration capability of a graphing utility, you obtain

$$A \approx 76,897.5 \text{ ft}^2.$$




Section 4.3 Riemann Sums and Definite Integrals

1. $f(x) = \sqrt{x}$, $y = 0$, $x = 0$, $x = 3$, $c_i = \frac{3i^2}{n^2}$

$$\Delta x_i = \frac{3i^2}{n^2} - \frac{3(i-1)^2}{n^2} = \frac{3}{n^2}(2i-1)$$

$$\begin{aligned} \lim_{n \rightarrow \infty} \sum_{i=1}^n f(c_i) \Delta x_i &= \lim_{n \rightarrow \infty} \sum_{i=1}^n \sqrt{\frac{3i^2}{n^2}} \frac{3}{n^2}(2i-1) \\ &= \lim_{n \rightarrow \infty} \frac{3\sqrt{3}}{n^3} \sum_{i=1}^n (2i^2 - i) \\ &= \lim_{n \rightarrow \infty} \frac{3\sqrt{3}}{n^3} \left[2 \frac{n(n+1)(2n+1)}{6} - \frac{n(n+1)}{2} \right] \\ &= \lim_{n \rightarrow \infty} 3\sqrt{3} \left[\frac{(n+1)(2n+1)}{3n^2} - \frac{n+1}{2n^2} \right] \\ &= 3\sqrt{3} \left[\frac{2}{3} - 0 \right] = 2\sqrt{3} \approx 3.464 \end{aligned}$$



3. $y = 6$ on $[4, 10]$. **Note:** $\Delta x = \frac{10 - 4}{n} = \frac{6}{n}$, $\|\Delta\| \rightarrow 0$ as $n \rightarrow \infty$

$$\sum_{i=1}^n f(c_i) \Delta x_i = \sum_{i=1}^n f\left(4 + \frac{6i}{n}\right)\left(\frac{6}{n}\right) = \sum_{i=1}^n 6\left(\frac{6}{n}\right) = \sum_{i=1}^n \frac{36}{n} = 36$$

$$\int_4^{10} 6 \, dx = \lim_{n \rightarrow \infty} 36 = 36$$

5. $y = x^3$ on $[-1, 1]$. **Note:** $\Delta x = \frac{1 - (-1)}{n} = \frac{2}{n}$, $\|\Delta\| \rightarrow 0$ as $n \rightarrow \infty$

$$\begin{aligned} \sum_{i=1}^n f(c_i) \Delta x_i &= \sum_{i=1}^n f\left(-1 + \frac{2i}{n}\right)\left(\frac{2}{n}\right) = \sum_{i=1}^n \left(-1 + \frac{2i}{n}\right)^3 \left(\frac{2}{n}\right) = \sum_{i=1}^n \left[-1 + \frac{6i}{n} - \frac{12i^2}{n^2} + \frac{8i^3}{n^3}\right] \left(\frac{2}{n}\right) \\ &= -2 + \frac{12}{n^2} \sum_{i=1}^n i - \frac{24}{n^3} \sum_{i=1}^n i^2 + \frac{16}{n^4} \sum_{i=1}^n i^3 \\ &= -2 + 6\left(1 + \frac{1}{n}\right) - 4\left(2 + \frac{3}{n} + \frac{1}{n^2}\right) + 4\left(1 + \frac{2}{n} + \frac{1}{n^2}\right) = \frac{2}{n} \end{aligned}$$

$$\int_{-1}^1 x^3 \, dx = \lim_{n \rightarrow \infty} \frac{2}{n} = 0$$

7. $y = x^2 + 1$ on $[1, 2]$. **Note:** $\Delta x = \frac{2 - 1}{n} = \frac{1}{n}$, $\|\Delta\| \rightarrow 0$ as $n \rightarrow \infty$

$$\begin{aligned} \sum_{i=1}^n f(c_i) \Delta x_i &= \sum_{i=1}^n f\left(1 + \frac{i}{n}\right)\left(\frac{1}{n}\right) = \sum_{i=1}^n \left[\left(1 + \frac{i}{n}\right)^2 + 1\right] \left(\frac{1}{n}\right) = \sum_{i=1}^n \left[1 + \frac{2i}{n} + \frac{i^2}{n^2} + 1\right] \left(\frac{1}{n}\right) \\ &= 2 + \frac{2}{n^2} \sum_{i=1}^n i + \frac{1}{n^3} \sum_{i=1}^n i^2 = 2 + \left(1 + \frac{1}{n}\right) + \frac{1}{6} \left(2 + \frac{3}{n} + \frac{1}{n^2}\right) = \frac{10}{3} + \frac{3}{2n} + \frac{1}{6n^2} \end{aligned}$$

$$\int_1^2 (x^2 + 1) \, dx = \lim_{n \rightarrow \infty} \left(\frac{10}{3} + \frac{3}{2n} + \frac{1}{6n^2}\right) = \frac{10}{3}$$

9. $\lim_{\|\Delta\| \rightarrow 0} \sum_{i=1}^n (3c_i + 10) \Delta x_i = \int_{-1}^5 (3x + 10) \, dx$

on the interval $[-1, 5]$.

11. $\lim_{\|\Delta\| \rightarrow 0} \sum_{i=1}^n \sqrt{c_i^2 + 4} \Delta x_i = \int_0^3 \sqrt{x^2 + 4} \, dx$

on the interval $[0, 3]$.

13. $\int_0^5 3 \, dx$

15. $\int_{-4}^4 (4 - |x|) \, dx$

17. $\int_{-2}^2 (4 - x^2) \, dx$


19. $\int_0^\pi \sin x \, dx$

21. $\int_0^2 y^3 \, dy$

23. Rectangle

$$A = bh = 3(4)$$


$$A = \int_0^3 4 \, dx = 12$$



25. Triangle


$$A = \frac{1}{2}bh = \frac{1}{2}(4)(4)$$

$$A = \int_0^4 x \, dx = 8$$

**29. Triangle**

$$A = \frac{1}{2}bh = \frac{1}{2}(2)(1)$$

$$A = \int_{-1}^1 (1 - |x|) \, dx = 1$$



In Exercises 33–39, $\int_2^4 x^3 \, dx = 60$, $\int_2^4 x \, dx = 6$, $\int_2^4 1 \, dx = 2$

$$33. \int_4^2 x \, dx = - \int_2^4 x \, dx = -6$$

$$37. \int_2^4 (x - 8) \, dx = \int_2^4 x \, dx - 8 \int_2^4 1 \, dx = 6 - 8(2) = -10$$

$$41. (a) \int_0^7 f(x) \, dx = \int_0^5 f(x) \, dx + \int_5^7 f(x) \, dx = 10 + 3 = 13$$

$$(b) \int_5^0 f(x) \, dx = - \int_0^5 f(x) \, dx = -10$$

$$(c) \int_5^5 f(x) \, dx = 0$$

$$(d) \int_0^5 3f(x) \, dx = 3 \int_0^5 f(x) \, dx = 3(10) = 30$$

$$45. (a) \text{Quarter circle below } x\text{-axis: } -\frac{1}{4}\pi r^2 = -\frac{1}{4}\pi(2)^2 = -\pi$$

$$(b) \text{Triangle: } \frac{1}{2}bh = \frac{1}{2}(4)(2) = 4$$

$$(c) \text{Triangle + Semicircle below } x\text{-axis: } -\frac{1}{2}(2)(1) - \frac{1}{2}\pi(2)^2 = -(1 + 2\pi)$$

$$(d) \text{Sum of parts (b) and (c): } 4 - (1 + 2\pi) = 3 - 2\pi$$

$$(e) \text{Sum of absolute values of (b) and (c): } 4 + (1 + 2\pi) = 5 + 2\pi$$


$$(f) \text{Answer to (d) plus } 2(10) = 20: (3 - 2\pi) + 20 = 23 - 2\pi$$

47. The left endpoint approximation will be greater than the actual area: $>$

27. Trapezoid


$$A = \frac{b_1 + b_2}{2}h = \left(\frac{5+9}{2}\right)2$$

$$A = \int_0^2 (2x + 5) \, dx = 14$$

**31. Semicircle**

$$A = \frac{1}{2}\pi r^2 = \frac{1}{2}\pi(3)^2$$

$$A = \int_{-3}^3 \sqrt{9 - x^2} \, dx = \frac{9\pi}{2}$$



$$35. \int_2^4 4x \, dx = 4 \int_2^4 x \, dx = 4(6) = 24$$

$$39. \int_2^4 \left(\frac{1}{2}x^3 - 3x + 2 \right) \, dx = \frac{1}{2} \int_2^4 x^3 \, dx - 3 \int_2^4 x \, dx + 2 \int_2^4 1 \, dx \\ = \frac{1}{2}(60) - 3(6) + 2(2) = 16$$

$$43. (a) \int_2^6 [f(x) + g(x)] \, dx = \int_2^6 f(x) \, dx + \int_2^6 g(x) \, dx \\ = 10 + (-2) = 8$$

$$(b) \int_2^6 [g(x) - f(x)] \, dx = \int_2^6 g(x) \, dx - \int_2^6 f(x) \, dx \\ = -2 - 10 = -12$$


$$(c) \int_2^6 2g(x) \, dx = 2 \int_2^6 g(x) \, dx = 2(-2) = -4$$

$$(d) \int_2^6 3f(x) \, dx = 3 \int_2^6 f(x) \, dx = 3(10) = 30$$

49. Because the curve is concave upward, the midpoint approximation will be less than the actual area: $<$


51. $f(x) = \frac{1}{x-4}$

is not integrable on the interval $[3, 5]$ and f has a discontinuity at $x = 4$.



a. $A \approx 5$ square units

55.



d. $\int_0^1 2 \sin \pi x \, dx \approx \frac{1}{2}(1)(2) \approx 1$

57. $\int_0^3 x \sqrt{3-x} \, dx$

n	4	8	12	16	20
$L(n)$	3.6830	3.9956	4.0707	4.1016	4.1177
$M(n)$	4.3082	4.2076	4.1838	4.1740	4.1690
$R(n)$	3.6830	3.9956	4.0707	4.1016	4.1177

59. $\int_0^{\pi/2} \sin^2 x \, dx$

n	4	8	12	16	20
$L(n)$	0.5890	0.6872	0.7199	0.7363	0.7461
$M(n)$	0.7854	0.7854	0.7854	0.7854	0.7854
$R(n)$	0.9817	0.8836	0.8508	0.8345	0.8247

61. True

63. True

65. False

$$\int_0^2 (-x) \, dx = -2$$

67. $f(x) = x^2 + 3x, [0, 8]$

$x_0 = 0, x_1 = 1, x_2 = 3, x_3 = 7, x_4 = 8$

$\Delta x_1 = 1, \Delta x_2 = 2, \Delta x_3 = 4, \Delta x_4 = 1$

$c_1 = 1, c_2 = 2, c_3 = 5, c_4 = 8$

$$\sum_{i=1}^4 f(c_i) \Delta x = f(1) \Delta x_1 + f(2) \Delta x_2 + f(5) \Delta x_3 + f(8) \Delta x_4$$

$$= (4)(1) + (10)(2) + (40)(4) + (88)(1) = 272$$

69. $f(x) = \begin{cases} 1, & x \text{ is rational} \\ 0, & x \text{ is irrational} \end{cases}$

is not integrable on the interval $[0, 1]$. As $\|\Delta\| \rightarrow 0$, $f(c_i) = 1$ or $f(c_i) = 0$ in each subinterval since there are an infinite number of both rational and irrational numbers in any interval, no matter how small.


71. Let $f(x) = x^2$, $0 \leq x \leq 1$, and $\Delta x_i = 1/n$. The appropriate Riemann Sum is

$$\begin{aligned}\sum_{i=1}^n f(c_i)\Delta x_i &= \sum_{i=1}^n \left(\frac{i}{n}\right)^2 \frac{1}{n} = \frac{1}{n^3} \sum_{i=1}^n i^2. \\ \lim_{n \rightarrow \infty} \frac{1}{n^3} [1^2 + 2^2 + 3^2 + \dots + n^2] &= \lim_{n \rightarrow \infty} \frac{1}{n^3} \cdot \frac{n(2n+1)(n+1)}{6} \\ &= \lim_{n \rightarrow \infty} \frac{2n^2 + 3n + 1}{6n^2} = \lim_{n \rightarrow \infty} \left(\frac{1}{3} + \frac{1}{2n} + \frac{1}{6n^2}\right) = \frac{1}{3}\end{aligned}$$

Section 4.4 The Fundamental Theorem of Calculus


1. $f(x) = \frac{4}{x^2 + 1}$

$\int_0^\pi \frac{4}{x^2 + 1} dx$ is positive.



3. $f(x) = x\sqrt{x^2 + 1}$

$$\int_{-2}^2 x\sqrt{x^2 + 1} dx = 0$$



5. $\int_0^1 2x dx = \left[x^2 \right]_0^1 = 1 - 0 = 1$

7. $\int_{-1}^0 (x - 2) dx = \left[\frac{x^2}{2} - 2x \right]_{-1}^0 = 0 - \left(\frac{1}{2} + 2 \right) = -\frac{5}{2}$

9. $\int_{-1}^1 (t^2 - 2) dt = \left[\frac{t^3}{3} - 2t \right]_{-1}^1 = \left(\frac{1}{3} - 2 \right) - \left(-\frac{1}{3} + 2 \right) = -\frac{10}{3}$

11. $\int_0^1 (2t - 1)^2 dt = \int_0^1 (4t^2 - 4t + 1) dt = \left[\frac{4}{3}t^3 - 2t^2 + t \right]_0^1 = \frac{4}{3} - 2 + 1 = \frac{1}{3}$

13. $\int_1^2 \left(\frac{3}{x^2} - 1 \right) dx = \left[-\frac{3}{x} - x \right]_1^2 = \left(-\frac{3}{2} - 2 \right) - (-3 - 1) = \frac{1}{2}$

15. $\int_1^4 \frac{u-2}{\sqrt{u}} du = \int_1^4 (u^{1/2} - 2u^{-1/2}) du = \left[\frac{2}{3}u^{3/2} - 4u^{1/2} \right]_1^4 = \left[\frac{2}{3}(\sqrt{4})^3 - 4\sqrt{4} \right] - \left[\frac{2}{3} - 4 \right] = \frac{2}{3}$

17. $\int_{-1}^1 (\sqrt[3]{t} - 2) dt = \left[\frac{3}{4}t^{4/3} - 2t \right]_{-1}^1 = \left(\frac{3}{4} - 2 \right) - \left(\frac{3}{4} + 2 \right) = -4$

19. $\int_0^1 \frac{x - \sqrt{x}}{3} dx = \frac{1}{3} \int_0^1 (x - x^{1/2}) dx = \frac{1}{3} \left[\frac{x^2}{2} - \frac{2}{3}x^{3/2} \right]_0^1 = \frac{1}{3} \left(\frac{1}{2} - \frac{2}{3} \right) = -\frac{1}{18}$

21. $\int_{-1}^0 (t^{1/3} - t^{2/3}) dt = \left[\frac{3}{4}t^{4/3} - \frac{3}{5}t^{5/3} \right]_{-1}^0 = 0 - \left(\frac{3}{4} + \frac{3}{5} \right) = -\frac{27}{20}$

23. $\int_0^3 |2x - 3| dx = \int_0^{3/2} (3 - 2x) dx + \int_{3/2}^3 (2x - 3) dx \quad \left(\text{split up the integral at the zero } x = \frac{3}{2} \right)$

$$= \left[3x - x^2 \right]_0^{3/2} + \left[x^2 - 3x \right]_{3/2}^3 = \left(\frac{9}{2} - \frac{9}{4} \right) - 0 + (9 - 9) - \left(\frac{9}{4} - \frac{9}{2} \right) = 2 \left(\frac{9}{2} - \frac{9}{4} \right) = \frac{9}{2}$$

$$\begin{aligned}
 25. \int_0^3 |x^2 - 4| dx &= \int_0^2 (4 - x^2) dx + \int_2^3 (x^2 - 4) dx \\
 &= \left[4x - \frac{x^3}{3} \right]_0^2 + \left[\frac{x^3}{3} - 4x \right]_2^3 \\
 &= \left(8 - \frac{8}{3} \right) + (9 - 12) - \left(\frac{8}{3} - 8 \right) \\
 &= \frac{23}{3}
 \end{aligned}$$

$$27. \int_0^\pi (1 + \sin x) dx = \left[x - \cos x \right]_0^\pi = (\pi + 1) - (0 - 1) = 2 + \pi$$

$$29. \int_{-\pi/6}^{\pi/6} \sec^2 x dx = \left[\tan x \right]_{-\pi/6}^{\pi/6} = \frac{\sqrt{3}}{3} - \left(-\frac{\sqrt{3}}{3} \right) = \frac{2\sqrt{3}}{3}$$

$$31. \int_{-\pi/3}^{\pi/3} 4 \sec \theta \tan \theta d\theta = \left[4 \sec \theta \right]_{-\pi/3}^{\pi/3} = 4(2) - 4(2) = 0$$

$$33. \int_0^3 10,000(t - 6) dt = 10,000 \left[\frac{t^2}{2} - 6t \right]_0^3 = -\$135,000$$

$$35. A = \int_0^1 (x - x^2) dx = \left[\frac{x^2}{2} - \frac{x^3}{3} \right]_0^1 = \frac{1}{6}$$

$$37. A = \int_0^3 (3 - x)\sqrt{x} dx = \int_0^3 (3x^{1/2} - x^{3/2}) dx = \left[2x^{3/2} - \frac{2}{5}x^{5/2} \right]_0^3 = \left[\frac{x\sqrt{x}}{5}(10 - 2x) \right]_0^3 = \frac{12\sqrt{3}}{5}$$

$$39. A = \int_0^{\pi/2} \cos x dx = \left[\sin x \right]_0^{\pi/2} = 1$$

41. Since $y = 0$ on $[0, 2]$,

$$A = \int_0^2 (3x^2 + 1) dx = \left[x^3 + x \right]_0^2 = 8 + 2 = 10.$$

43. Since $y = 0$ on $[0, 2]$,

$$A = \int_0^2 (x^3 + x) dx = \left[\frac{x^4}{4} + \frac{x^2}{2} \right]_0^2 = 4 + 2 = 6.$$

$$45. \int_0^2 (x - 2\sqrt{x}) dx = \left[\frac{x^2}{2} - \frac{4x^{3/2}}{3} \right]_0^2 = 2 - \frac{8\sqrt{2}}{3}$$

$$f(c)(2 - 0) = \frac{6 - 8\sqrt{2}}{3}$$

$$c - 2\sqrt{c} = \frac{3 - 4\sqrt{2}}{3}$$

$$c - 2\sqrt{c} + 1 = \frac{3 - 4\sqrt{2}}{3} + 1$$

$$(\sqrt{c} - 1)^2 = \frac{6 - 4\sqrt{2}}{3}$$

$$\sqrt{c} - 1 = \pm \sqrt{\frac{6 - 4\sqrt{2}}{3}}$$

$$c = \left[1 \pm \sqrt{\frac{6 - 4\sqrt{2}}{3}} \right]^2$$

$$c \approx 0.4380 \text{ or } c \approx 1.7908$$

47. $\int_{-\pi/4}^{\pi/4} 2 \sec^2 x \, dx = \left[2 \tan x \right]_{-\pi/4}^{\pi/4} = 2(1) - 2(-1) = 4$

$$f(c) \left[\frac{\pi}{4} - \left(-\frac{\pi}{4} \right) \right] = 4$$

$$2 \sec^2 c = \frac{8}{\pi}$$

$$\sec^2 c = \frac{4}{\pi}$$

$$\sec c = \pm \frac{2}{\sqrt{\pi}}$$


$$c = \pm \text{arcsec} \left(\frac{2}{\sqrt{\pi}} \right)$$

$$= \pm \arccos \frac{\sqrt{\pi}}{2} \approx \pm 0.4817$$

49. $\frac{1}{2 - (-2)} \int_{-2}^2 (4 - x^2) \, dx = \frac{1}{4} \left[4x - \frac{1}{3}x^3 \right]_{-2}^2 = \frac{1}{4} \left[\left(8 - \frac{8}{3} \right) - \left(-8 + \frac{8}{3} \right) \right] = \frac{8}{3}$

$$\text{Average value} = \frac{8}{3}$$

$$4 - x^2 = \frac{8}{3} \text{ when } x^2 = 4 - \frac{8}{3} \text{ or } x = \pm \frac{2\sqrt{3}}{3} \approx \pm 1.155.$$




51. $\frac{1}{\pi - 0} \int_0^\pi \sin x \, dx = \left[-\frac{1}{\pi} \cos x \right]_0^\pi = \frac{2}{\pi}$

$$\text{Average value} = \frac{2}{\pi}$$

$$\sin x = \frac{2}{\pi}$$

$$x \approx 0.690, 2.451$$



55. $\int_0^2 f(x) \, dx = -(\text{area of region A}) = -1.5$

53. If f is continuous on $[a, b]$ and $F'(x) = f(x)$ on $[a, b]$, then $\int_a^b f(x) \, dx = F(b) - F(a)$.

57. $\int_0^6 |f(x)| \, dx = -\int_0^2 f(x) \, dx + \int_2^6 f(x) \, dx = 1.5 + 5.0 = 6.5$

59. $\int_0^6 [2 + f(x)] \, dx = \int_0^6 2 \, dx + \int_0^6 f(x) \, dx$
 $= 12 + 3.5 = 15.5$


61. (a) $F(x) = k \sec^2 x$

$$F(0) = k = 500$$


$$F(x) = 500 \sec^2 x$$

(b) $\frac{1}{\pi/3 - 0} \int_0^{\pi/3} 500 \sec^2 x \, dx = \frac{1500}{\pi} \left[\tan x \right]_0^{\pi/3}$
 $= \frac{1500}{\pi} (\sqrt{3} - 0)$
 $\approx 826.99 \text{ newtons}$
 $\approx 827 \text{ newtons}$

63. $\frac{1}{5 - 0} \int_0^5 (0.1729t + 0.1552t^2 - 0.0374t^3) \, dt \approx \frac{1}{5} \left[0.08645t^2 + 0.05073t^3 - 0.00935t^4 \right]_0^5 \approx 0.5318 \text{ liter}$




The area above the x -axis equals the area below the x -axis. Thus, the average value is zero.



The average value of S appears to be g .

67. (a) $v = -8.61 \times 10^{-4}t^3 + 0.0782t^2 - 0.208t + 0.0952$



(c) $\int_0^{60} v(t) dt = \left[\frac{-8.61 \times 10^{-4}t^4}{4} + \frac{0.0782t^3}{3} - \frac{0.208t^2}{2} + 0.0952t \right]_0^{60} \approx 2476 \text{ meters}$

69. $F(x) = \int_0^x (t - 5) dt = \left[\frac{t^2}{2} - 5t \right]_0^x = \frac{x^2}{2} - 5x$

$$F(2) = \frac{4}{2} - 5(2) = -8$$

$$F(5) = \frac{25}{2} - 5(5) = -\frac{25}{2}$$

$$F(8) = \frac{64}{2} - 5(8) = -8$$

71. $F(x) = \int_1^x \frac{10}{v^2} dv = \int_1^x 10v^{-2} dv = \frac{-10}{v} \Big|_1^x$

$$F(2) = 10 \left(\frac{1}{2} \right) = 5$$

$$\begin{aligned} F(5) &= 10 \left(\frac{4}{5} \right) = 8 \\ F(8) &= 10 \left(\frac{7}{8} \right) = \frac{35}{4} \end{aligned}$$

73. $F(x) = \int_1^x \cos \theta d\theta = \sin \theta \Big|_1^x = \sin x - \sin 1$

$$F(2) = \sin 2 - \sin 1 = 0.0678$$

$$F(5) = \sin 5 - \sin 1 \approx -1.8004$$

$$F(8) = \sin 8 - \sin 1 \approx 0.1479$$

75. (a) $\int_0^x (t + 2) dt = \left[\frac{t^2}{2} + 2t \right]_0^x = \frac{1}{2}x^2 + 2x$

(b) $\frac{d}{dx} \left[\frac{1}{2}x^2 + 2x \right] = x + 2$

77. (a) $\int_8^x \sqrt[3]{t} dt = \left[\frac{3}{4}t^{4/3} \right]_8^x = \frac{3}{4}(x^{4/3} - 16) = \frac{3}{4}x^{4/3} - 12$

(b) $\frac{d}{dx} \left[\frac{3}{4}x^{4/3} - 12 \right] = x^{1/3} = \sqrt[3]{x}$

81. $F(x) = \int_{-2}^x (t^2 - 2t) dt$

$$F'(x) = x^2 - 2x$$

83. $F(x) = \int_{-1}^x \sqrt{t^4 + 1} dt$

$$F'(x) = \sqrt{x^4 + 1}$$

79. (a) $\int_{x/4}^x \sec^2 t dt = \left[\tan t \right]_{x/4}^x = \tan x - 1$

(b) $\frac{d}{dx} [\tan x - 1] = \sec^2 x$

85. $F(x) = \int_0^x t \cos t dt$

$$F'(x) = x \cos x$$


$$\begin{aligned}
 87. \quad F(x) &= \int_x^{x+2} (4t + 1) dt \\
 &= \left[2t^2 + t \right]_x^{x+2} \\
 &= [2(x+2)^2 + (x+2)] - [2x^2 + x] \\
 &= 8x + 10 \\
 F'(x) &= 8
 \end{aligned}$$

$$\begin{aligned}
 89. \quad F(x) &= \int_0^{\sin x} \sqrt{t} dt = \left[\frac{2}{3} t^{3/2} \right]_0^{\sin x} = \frac{2}{3} (\sin x)^{3/2} \\
 F'(x) &= (\sin x)^{1/2} \cos x = \cos x \sqrt{\sin x}
 \end{aligned}$$

Alternate solution

$$\begin{aligned}
 F(x) &= \int_0^{\sin x} \sqrt{t} dt \\
 F'(x) &= \sqrt{\sin x} \frac{d}{dx} (\sin x) = \sqrt{\sin x} (\cos x)
 \end{aligned}$$

$$\begin{aligned}
 93. \quad g(x) &= \int_0^x f(t) dt \\
 g(0) &= 0, g(1) \approx \frac{1}{2}, g(2) \approx 1, g(3) \approx \frac{1}{2}, g(4) = 0
 \end{aligned}$$



g has a relative maximum at $x = 2$.

97. True

$$101. \quad f(x) = \int_0^{1/x} \frac{1}{t^2 + 1} dt + \int_0^x \frac{1}{t^2 + 1} dt$$

By the Second Fundamental Theorem of Calculus, we have

$$\begin{aligned}
 f'(x) &= \frac{1}{(1/x)^2 + 1} \left(-\frac{1}{x^2} \right) + \frac{1}{x^2 + 1} \\
 &= -\frac{1}{1+x^2} + \frac{1}{x^2+1} = 0.
 \end{aligned}$$

Since $f'(x) = 0$, $f(x)$ must be constant.

Alternate solution:

$$\begin{aligned}
 F(x) &= \int_x^{x+2} (4t + 1) dt \\
 &= \int_x^0 (4t + 1) dt + \int_0^{x+2} (4t + 1) dt \\
 &= -\int_0^x (4t + 1) dt + \int_0^{x+2} (4t + 1) dt \\
 F'(x) &= -(4x + 1) + 4(x + 2) + 1 = 8
 \end{aligned}$$

$$\begin{aligned}
 91. \quad F(x) &= \int_0^{x^3} \sin t^2 dt \\
 F'(x) &= \sin(x^3)^2 \cdot 3x^2 = 3x^2 \sin x^6
 \end{aligned}$$

$$\begin{aligned}
 95. \quad (a) \quad C(x) &= 5000 \left(25 + 3 \int_0^x t^{1/4} dt \right) \\
 &= 5000 \left(25 + 3 \left[\frac{4}{5} t^{5/4} \right]_0^x \right) \\
 &= 5000 \left(25 + \frac{12}{5} x^{5/4} \right) = 1000(125 + 12x^{5/4}) \\
 (b) \quad C(1) &= 1000(125 + 12(1)) = \$137,000 \\
 C(5) &= 1000(125 + 12(5)^{5/4}) \approx \$214,721 \\
 C(10) &= 1000(125 + 12(10)^{5/4}) \approx \$338,394
 \end{aligned}$$

$$99. \quad \text{False;} \quad \int_{-1}^1 x^{-2} dx = \int_{-1}^0 x^{-2} dx + \int_0^1 x^{-2} dx$$

Each of these integrals is infinite. $f(x) = x^{-2}$ has a nonremovable discontinuity at $x = 0$.

103. $x(t) = t^3 - 6t^2 + 9t - 2$

$$x'(t) = 3t^2 - 12t + 9$$

$$= 3(t^2 - 4t + 3)$$

$$= 3(t - 3)(t - 1)$$

$$\begin{aligned} \text{Total distance} &= \int_0^5 |x'(t)| dt \\ &= \int_0^5 3|(t - 3)(t - 1)| dt \\ &= 3 \int_0^1 (t^2 - 4t + 3) dt - 3 \int_1^3 (t^2 - 4t + 3) dt + 3 \int_3^5 (t^2 - 4t + 3) dt \\ &= 4 + 4 + 20 \\ &= 28 \text{ units} \end{aligned}$$

105. Total distance = $\int_1^4 |x'(t)| dt$

$$= \int_1^4 |v(t)| dt$$

$$= \int_1^4 \frac{1}{\sqrt{t}} dt$$

$$= 2t^{1/2} \Big|_1^4$$

$$= 2(2 - 1) = 2 \text{ units}$$

Section 4.5 Integration by Substitution

$$\underline{\int f(g(x))g'(x) dx} \quad \underline{u = g(x)} \quad \underline{du = g'(x) dx}$$

1. $\int (5x^2 + 1)^2(10x) dx$ $5x^2 + 1$ $10x dx$

3. $\int \frac{x}{\sqrt{x^2 + 1}} dx$ $x^2 + 1$ $2x dx$

5. $\int \tan^2 x \sec^2 x dx$ $\tan x$ $\sec^2 x dx$

7. $\int (1 + 2x)^4 2 dx = \frac{(1 + 2x)^5}{5} + C$

Check: $\frac{d}{dx} \left[\frac{(1 + 2x)^5}{5} + C \right] = 2(1 + 2x)^4$

9. $\int (9 - x^2)^{1/2}(-2x) dx = \frac{(9 - x^2)^{3/2}}{3/2} + C = \frac{2}{3}(9 - x^2)^{3/2} + C$

Check: $\frac{d}{dx} \left[\frac{2}{3}(9 - x^2)^{3/2} + C \right] = \frac{2}{3} \cdot \frac{3}{2}(9 - x^2)^{1/2}(-2x) = \sqrt{9 - x^2}(-2x)$

11. $\int x^3(x^4 + 3)^2 dx = \frac{1}{4} \int (x^4 + 3)^2(4x^3) dx = \frac{1}{4} \frac{(x^4 + 3)^3}{3} + C = \frac{(x^4 + 3)^3}{12} + C$

Check: $\frac{d}{dx} \left[\frac{(x^4 + 3)^3}{12} + C \right] = \frac{3(x^4 + 3)^2}{12}(4x^3) = (x^4 + 3)^2(x^3)$

13. $\int x^2(x^3 - 1)^4 dx = \frac{1}{3} \int (x^3 - 1)^4(3x^2) dx = \frac{1}{3} \left[\frac{(x^3 - 1)^5}{5} \right] + C = \frac{(x^3 - 1)^5}{15} + C$

Check: $\frac{d}{dx} \left[\frac{(x^3 - 1)^5}{15} + C \right] = \frac{5(x^3 - 1)^4(3x^2)}{15} = x^2(x^3 - 1)^4$

15. $\int t\sqrt{t^2 + 2} dt = \frac{1}{2} \int (t^2 + 2)^{1/2}(2t) dt = \frac{1}{2} \frac{(t^2 + 2)^{3/2}}{3/2} + C = \frac{(t^2 + 2)^{3/2}}{3} + C$

Check: $\frac{d}{dt} \left[\frac{(t^2 + 2)^{3/2}}{3} + C \right] = \frac{3/2(t^2 + 2)^{1/2}(2t)}{3} = (t^2 + 2)^{1/2}t$

17. $\int 5x(1 - x^2)^{1/3} dx = -\frac{5}{2} \int (1 - x^2)^{1/3}(-2x) dx = -\frac{5}{2} \cdot \frac{(1 - x^2)^{4/3}}{4/3} + C = -\frac{15}{8}(1 - x^2)^{4/3} + C$

Check: $\frac{d}{dx} \left[-\frac{15}{8}(1 - x^2)^{4/3} + C \right] = -\frac{15}{8} \cdot \frac{4}{3}(1 - x^2)^{1/3}(-2x) = 5x(1 - x^2)^{1/3} = 5x\sqrt[3]{1 - x^2}$

19. $\int \frac{x}{(1 - x^2)^3} dx = -\frac{1}{2} \int (1 - x^2)^{-3}(-2x) dx = -\frac{1}{2} \frac{(1 - x^2)^{-2}}{-2} + C = \frac{1}{4(1 - x^2)^2} + C$

Check: $\frac{d}{dx} \left[\frac{1}{4(1 - x^2)^2} + C \right] = \frac{1}{4}(-2)(1 - x^2)^{-3}(-2x) = \frac{x}{(1 - x^2)^3}$

21. $\int \frac{x^2}{(1 + x^3)^2} dx = \frac{1}{3} \int (1 + x^3)^{-2}(3x^2) dx = \frac{1}{3} \left[\frac{(1 + x^3)^{-1}}{-1} \right] + C = -\frac{1}{3(1 + x^3)} + C$

Check: $\frac{d}{dx} \left[-\frac{1}{3(1 + x^3)} + C \right] = -\frac{1}{3}(-1)(1 + x^3)^{-2}(3x^2) = \frac{x^2}{(1 + x^3)^2}$

23. $\int \frac{x}{\sqrt{1 - x^2}} dx = -\frac{1}{2} \int (1 - x^2)^{-1/2}(-2x) dx = -\frac{1}{2} \frac{(1 - x^2)^{1/2}}{1/2} + C = -\sqrt{1 - x^2} + C$

Check: $\frac{d}{dx} \left[-\frac{1}{2}(1 - x^2)^{-1/2} + C \right] = -\frac{1}{2}(1 - x^2)^{-1/2}(-2x) = \frac{x}{\sqrt{1 - x^2}}$

25. $\int \left(1 + \frac{1}{t}\right)^3 \left(\frac{1}{t^2}\right) dt = - \int \left(1 + \frac{1}{t}\right)^3 \left(-\frac{1}{t^2}\right) dt = -\frac{[1 + (1/t)]^4}{4} + C$

Check: $\frac{d}{dt} \left[-\frac{[1 + (1/t)]^4}{4} + C \right] = -\frac{1}{4}(4)\left(1 + \frac{1}{t}\right)^3 \left(-\frac{1}{t^2}\right) = \frac{1}{t^2} \left(1 + \frac{1}{t}\right)^3$

27. $\int \frac{1}{\sqrt{2x}} dx = \frac{1}{2} \int (2x)^{-1/2} 2 dx = \frac{1}{2} \left[\frac{(2x)^{1/2}}{1/2} \right] + C = \sqrt{2x} + C$

Check: $\frac{d}{dx} \left[\sqrt{2x} + C \right] = \frac{1}{2}(2x)^{-1/2}(2) = \frac{1}{\sqrt{2x}}$

29. $\int \frac{x^2 + 3x + 7}{\sqrt{x}} dx = \int (x^{3/2} + 3x^{1/2} + 7x^{-1/2}) dx = \frac{2}{5}x^{5/2} + 2x^{3/2} + 14x^{1/2} + C = \frac{2}{5}\sqrt{x}(x^2 + 5x + 35) + C$

Check: $\frac{d}{dx} \left[\frac{2}{5}x^{5/2} + 2x^{3/2} + 14x^{1/2} + C \right] = \frac{x^2 + 3x + 7}{\sqrt{x}}$

31. $\int t^2 \left(t - \frac{2}{t} \right) dt = \int (t^3 - 2t) dt = \frac{1}{4}t^4 - t^2 + C$

Check: $\frac{d}{dt} \left[\frac{1}{4}t^4 - t^2 + C \right] = t^3 - 2t = t^2 \left(t - \frac{2}{t} \right)$


33. $\int (9 - y)\sqrt{y} dy = \int (9y^{1/2} - y^{3/2}) dy = 9 \left(\frac{2}{3}y^{3/2} \right) - \frac{2}{5}y^{5/2} + C = \frac{2}{5}y^{3/2}(15 - y) + C$

Check: $\frac{d}{dy} \left[\frac{2}{5}y^{3/2}(15 - y) + C \right] = \frac{d}{dy} \left[6y^{3/2} - \frac{2}{5}y^{5/2} + C \right] = 9y^{1/2} - y^{3/2} = (9 - y)\sqrt{y}$

35. $y = \int \left[4x + \frac{4x}{\sqrt{16 - x^2}} \right] dx$
 $= 4 \int x dx - 2 \int (16 - x^2)^{-1/2}(-2x) dx$
 $= 4 \left(\frac{x^2}{2} \right) - 2 \left[\frac{(16 - x^2)^{1/2}}{1/2} \right] + C$
 $= 2x^2 - 4\sqrt{16 - x^2} + C$

37. $y = \int \frac{x+1}{(x^2 + 2x - 3)^2} dx$
 $= \frac{1}{2} \int (x^2 + 2x - 3)^{-2}(2x + 2) dx$
 $= \frac{1}{2} \left[\frac{(x^2 + 2x - 3)^{-1}}{-1} \right] + C$
 $= -\frac{1}{2(x^2 + 2x - 3)} + C$


39. (a)



(b) $\frac{dy}{dx} = x\sqrt{4 - x^2}, (2, 2)$

$$\begin{aligned} y &= \int x\sqrt{4 - x^2} dx = -\frac{1}{2} \int (4 - x^2)^{1/2}(-2x) dx \\ &= -\frac{1}{2} \cdot \frac{2}{3} (4 - x^2)^{3/2} + C = -\frac{1}{3}(4 - x^2)^{3/2} + C \\ (2, 2): 2 &= -\frac{1}{3}(4 - 2^2)^{3/2} + C \quad C = 2 \end{aligned}$$

$$y = -\frac{1}{3}(4 - x^2)^{3/2} + 2$$



41. $\int \pi \sin \pi x dx = -\cos \pi x + C$

43. $\int \sin 2x dx = \frac{1}{2} \int (\sin 2x)(2x) dx = -\frac{1}{2} \cos 2x + C$

45. $\int \frac{1}{\theta^2} \cos \frac{1}{\theta} d\theta = - \int \cos \frac{1}{\theta} \left(-\frac{1}{\theta^2} \right) d\theta = -\sin \frac{1}{\theta} + C$

47. $\int \sin 2x \cos 2x \, dx = \frac{1}{2} \int (\sin 2x)(2 \cos 2x) \, dx = \frac{1}{2} \frac{(\sin 2x)^2}{2} + C = \frac{1}{4} \sin^2 2x + C \quad \text{OR}$

$$\int \sin 2x \cos 2x \, dx = -\frac{1}{2} \int (\cos 2x)(-2 \sin 2x) \, dx = -\frac{1}{2} \frac{(\cos 2x)^2}{2} + C_1 = -\frac{1}{4} \cos^2 2x + C_1 \quad \text{OR}$$

$$\int \sin 2x \cos 2x \, dx = \frac{1}{2} \int 2 \sin 2x \cos 2x \, dx = \frac{1}{2} \int \sin 4x \, dx - \frac{1}{8} \cos 4x + C_2$$

49. $\int \tan^4 x \sec^2 x \, dx = \frac{\tan^5 x}{5} + C = \frac{1}{5} \tan^5 x + C$

51. $\int \frac{\csc^2 x}{\cot^3 x} \, dx = - \int (\cot x)^{-3} (-\csc^2 x) \, dx$
 $= -\frac{(\cot x)^{-2}}{-2} + C = \frac{1}{2 \cot^2 x} + C = \frac{1}{2} \tan^2 x + C = \frac{1}{2} (\sec^2 x - 1) + C = \frac{1}{2} \sec^2 x + C_1$

53. $\int \cot^2 x \, dx = \int (\csc^2 x - 1) \, dx = -\cot x - x + C$

55. $f(x) = \int \cos \frac{x}{2} \, dx = 2 \sin \frac{x}{2} + C$

Since $f(0) = 3 = 2 \sin 0 + C$, $C = 3$. Thus,

$$f(x) = 2 \sin \frac{x}{2} + 3.$$

57. $u = x + 2$, $x = u - 2$, $dx = du$

$$\begin{aligned} \int x \sqrt{x+2} \, dx &= \int (u-2)\sqrt{u} \, du \\ &= \int (u^{3/2} - 2u^{1/2}) \, du \\ &= \frac{2}{5}u^{5/2} - \frac{4}{3}u^{3/2} + C \\ &= \frac{2u^{3/2}}{15}(3u-10) + C \\ &= \frac{2}{15}(x+2)^{3/2}[3(x+2)-10] + C \\ &= \frac{2}{15}(x+2)^{3/2}(3x-4) + C \end{aligned}$$

59. $u = 1-x$, $x = 1-u$, $dx = -du$

$$\begin{aligned} \int x^2 \sqrt{1-x} \, dx &= - \int (1-u)^2 \sqrt{u} \, du \\ &= - \int (u^{1/2} - 2u^{3/2} + u^{5/2}) \, du \\ &= - \left(\frac{2}{3}u^{3/2} - \frac{4}{5}u^{5/2} + \frac{2}{7}u^{7/2} \right) + C \\ &= -\frac{2u^{3/2}}{105}(35-42u+15u^2) + C \\ &= -\frac{2}{105}(1-x)^{3/2}[35-42(1-x)+15(1-x)^2] + C \\ &= -\frac{2}{105}(1-x)^{3/2}(15x^2+12x+8) + C \end{aligned}$$

61. $u = 2x - 1, x = \frac{1}{2}(u + 1), dx = \frac{1}{2}du$

$$\begin{aligned}\int \frac{x^2 - 1}{\sqrt{2x - 1}} dx &= \int \frac{[(1/2)(u + 1)]^2 - 1}{\sqrt{u}} \frac{1}{2} du \\&= \frac{1}{8} \int u^{-1/2} [(u^2 + 2u + 1) - 4] du \\&= \frac{1}{8} \int (u^{3/2} + 2u^{1/2} - 3u^{-1/2}) du \\&= \frac{1}{8} \left(\frac{2}{5}u^{5/2} + \frac{4}{3}u^{3/2} - 6u^{1/2} \right) + C \\&= \frac{u^{1/2}}{60} (3u^2 + 10u - 45) + C \\&= \frac{\sqrt{2x - 1}}{60} [3(2x - 1)^2 + 10(2x - 1) - 45] + C \\&= \frac{1}{60} \sqrt{2x - 1} (12x^2 + 8x - 52) + C \\&= \frac{1}{15} \sqrt{2x - 1} (3x^2 + 2x - 13) + C\end{aligned}$$

63. $u = x + 1, x = u - 1, dx = du$

$$\begin{aligned}\int \frac{-x}{(x + 1) - \sqrt{x + 1}} dx &= \int \frac{-(u - 1)}{u - \sqrt{u}} du \\&= - \int \frac{(\sqrt{u} + 1)(\sqrt{u} - 1)}{\sqrt{u}(\sqrt{u} - 1)} du \\&= - \int (1 + u^{-1/2}) du \\&= -(u + 2u^{1/2}) + C \\&= -u - 2\sqrt{u} + C \\&= -(x + 1) - 2\sqrt{x + 1} + C \\&= -x - 2\sqrt{x + 1} - 1 + C \\&= -(x + 2\sqrt{x + 1}) + C_1\end{aligned}$$

where $C_1 = -1 + C$.

65. Let $u = x^2 + 1, du = 2x dx$.

$$\int_{-1}^1 x(x^2 + 1)^3 dx = \frac{1}{2} \int_{-1}^1 (x^2 + 1)^3 (2x) dx = \left[\frac{1}{8}(x^2 + 1)^4 \right]_{-1}^1 = 0$$

67. Let $u = x^3 + 1, du = 3x^2 dx$

$$\begin{aligned}\int_1^2 2x^2 \sqrt{x^3 + 1} dx &= 2 \cdot \frac{1}{3} \int_1^2 (x^3 + 1)^{1/2} (3x^2) dx \\&= \left[\frac{2}{3} \frac{(x^3 + 1)^{3/2}}{3/2} \right]_1^2 \\&= \frac{4}{9} \left[(x^3 + 1)^{3/2} \right]_1^2 \\&= \frac{4}{9} [27 - 2\sqrt{2}] = 12 - \frac{8}{9}\sqrt{2}\end{aligned}$$

69. Let $u = 2x + 1$, $du = 2 dx$.

$$\int_0^4 \frac{1}{\sqrt{2x+1}} dx = \frac{1}{2} \int_0^4 (2x+1)^{-1/2}(2) dx = \left[\sqrt{2x+1} \right]_0^4 = \sqrt{9} - \sqrt{1} = 2$$

71. Let $u = 1 + \sqrt{x}$, $du = \frac{1}{2\sqrt{x}} dx$.

$$\int_1^9 \frac{1}{\sqrt{x}(1+\sqrt{x})^2} dx = 2 \int_1^9 (1+\sqrt{x})^{-2} \left(\frac{1}{2\sqrt{x}} \right) dx = \left[-\frac{2}{1+\sqrt{x}} \right]_1^9 = -\frac{1}{2} + 1 = \frac{1}{2}$$

73. $u = 2 - x$, $x = 2 - u$, $dx = -du$

When $x = 1$, $u = 1$. When $x = 2$, $u = 0$.

$$\int_1^2 (x-1)\sqrt{2-x} dx = \int_1^0 -[(2-u)-1]\sqrt{u} du = \int_1^0 (u^{3/2} - u^{1/2}) du = \left[\frac{2}{5}u^{5/2} - \frac{2}{3}u^{3/2} \right]_1^0 = -\left[\frac{2}{5} - \frac{2}{3} \right] = \frac{4}{15}$$

75. $\int_0^{\pi/2} \cos\left(\frac{2}{3}x\right) dx = \left[\frac{3}{2} \sin\left(\frac{2}{3}x\right) \right]_0^{\pi/2} = \frac{3}{2} \left(\frac{\sqrt{3}}{2} \right) = \frac{3\sqrt{3}}{4}$

77. $u = x + 1$, $x = u - 1$, $dx = du$

When $x = 0$, $u = 1$. When $x = 7$, $u = 8$.

$$\begin{aligned} \text{Area} &= \int_0^7 x \sqrt[3]{x+1} dx = \int_1^8 (u-1) \sqrt[3]{u} du \\ &= \int_1^8 (u^{4/3} - u^{1/3}) du = \left[\frac{3}{7}u^{7/3} - \frac{3}{4}u^{4/3} \right]_1^8 = \left(\frac{384}{7} - 12 \right) - \left(\frac{3}{7} - \frac{3}{4} \right) = \frac{1209}{28} \end{aligned}$$




79. $A = \int_0^\pi (2 \sin x + \sin 2x) dx = -\left[2 \cos x + \frac{1}{2} \cos 2x \right]_0^\pi = 4$

81. Area $= \int_{\pi/2}^{2\pi/3} \sec^2\left(\frac{x}{2}\right) dx = 2 \int_{\pi/2}^{2\pi/3} \sec^2\left(\frac{x}{2}\right) \left(\frac{1}{2}\right) dx = \left[2 \tan\left(\frac{x}{2}\right) \right]_{\pi/2}^{2\pi/3} = 2(\sqrt{3} - 1)$

83. $\int_0^4 \frac{x}{\sqrt{2x+1}} dx \approx 3.333 = \frac{10}{3}$

85. $\int_3^7 x \sqrt{x-3} dx \approx 28.8 = \frac{144}{5}$

87. $\int_0^3 \left(\theta + \cos \frac{\theta}{6} \right) d\theta \approx 7.377$



89. $\int (2x-1)^2 dx = \frac{1}{2} \int (2x-1)^2 2 dx = \frac{1}{6} (2x-1)^3 + C_1 = \frac{4}{3}x^3 - 2x^2 + x - \frac{1}{6} + C_1$

$$\int (2x-1)^2 dx = \int (4x^2 - 4x + 1) dx = \frac{4}{3}x^3 - 2x^2 + x + C_2$$

They differ by a constant: $C_2 = C_1 - \frac{1}{6}$.

91. $f(x) = x^2(x^2 + 1)$ is even.

$$\begin{aligned}\int_{-2}^2 x^2(x^2 + 1) dx &= 2 \int_0^2 (x^4 + x^2) dx = 2 \left[\frac{x^5}{5} + \frac{x^3}{3} \right]_0^2 \\ &= 2 \left[\frac{32}{5} + \frac{8}{3} \right] = \frac{272}{15}\end{aligned}$$

93. $f(x) = x(x^2 + 1)^3$ is odd.

$$\int_{-2}^2 x(x^2 + 1)^3 dx = 0$$

95. $\int_0^2 x^2 dx = \left[\frac{x^3}{3} \right]_0^2 = \frac{8}{3}$; the function x^2 is an even function.

$$(a) \int_{-2}^0 x^2 dx = \int_0^2 x^2 dx = \frac{8}{3}$$

$$(b) \int_{-2}^2 x^2 dx = 2 \int_0^2 x^2 dx = \frac{16}{3}$$

$$(c) \int_0^2 (-x^2) dx = - \int_0^2 x^2 dx = -\frac{8}{3}$$

$$(d) \int_{-2}^0 3x^2 dx = 3 \int_0^2 x^2 dx = 8$$

$$\begin{aligned}\text{97. } \int_{-4}^4 (x^3 + 6x^2 - 2x - 3) dx &= \int_{-4}^4 (x^3 - 2x) dx + \int_{-4}^4 (6x^2 - 3) dx = 0 + 2 \int_0^4 (6x^2 - 3) dx = 2 \left[2x^3 - 3x \right]_0^4 = 232\end{aligned}$$

99. Answers will vary. See “Guidelines for Making a Change of Variables” on page 292.

101. $f(x) = x(x^2 + 1)^2$ is odd. Hence, $\int_{-2}^2 x(x^2 + 1)^2 dx = 0$.

$$\text{103. } \frac{dV}{dt} = \frac{k}{(t+1)^2}$$

$$V(t) = \int \frac{k}{(t+1)^2} dt = -\frac{k}{t+1} + C$$

$$V(0) = -k + C = 500,000$$

$$V(1) = -\frac{1}{2}k + C = 400,000$$

Solving this system yields $k = -200,000$ and $C = 300,000$. Thus,

$$V(t) = \frac{200,000}{t+1} + 300,000.$$

When $t = 4$, $V(4) = \$340,000$.

$$\text{105. } \frac{1}{b-a} \int_a^b \left[74.50 + 43.75 \sin \frac{\pi t}{6} \right] dt = \frac{1}{b-a} \left[74.50t - \frac{262.5}{\pi} \cos \frac{\pi t}{6} \right]_a^b$$

$$(a) \frac{1}{3} \left[74.50t - \frac{262.5}{\pi} \cos \frac{\pi t}{6} \right]_0^3 = \frac{1}{3} \left(223.5 + \frac{262.5}{\pi} \right) \approx 102.352 \text{ thousand units}$$

$$(b) \frac{1}{3} \left[74.50t - \frac{262.5}{\pi} \cos \frac{\pi t}{6} \right]_3^6 = \frac{1}{3} \left(447 + \frac{262.5}{\pi} - 223.5 \right) \approx 102.352 \text{ thousand units}$$

$$(c) \frac{1}{12} \left[74.50t - \frac{262.5}{\pi} \cos \frac{\pi t}{6} \right]_0^{12} = \frac{1}{12} \left(894 - \frac{262.5}{\pi} + \frac{262.5}{\pi} \right) = 74.5 \text{ thousand units}$$

107. $\frac{1}{b-a} \int_a^b [2 \sin(60\pi t) + \cos(120\pi t)] dt = \frac{1}{b-a} \left[-\frac{1}{30\pi} \cos(60\pi t) + \frac{1}{120\pi} \sin(120\pi t) \right]_a^b$

(a) $\frac{1}{(1/60)-0} \left[-\frac{1}{30\pi} \cos(60\pi t) + \frac{1}{120\pi} \sin(120\pi t) \right]_0^{1/60} = 60 \left[\left(\frac{1}{30\pi} + 0 \right) - \left(-\frac{1}{30\pi} \right) \right] = \frac{4}{\pi} \approx 1.273 \text{ amps}$

(b) $\frac{1}{(1/240)-0} \left[-\frac{1}{30\pi} \cos(60\pi t) + \frac{1}{120\pi} \sin(120\pi t) \right]_0^{1/240} = 240 \left[\left(-\frac{1}{30\sqrt{2}\pi} + \frac{1}{120\pi} \right) - \left(-\frac{1}{30\pi} \right) \right]$
 $= \frac{2}{\pi} (5 - 2\sqrt{2}) \approx 1.382 \text{ amps}$

(c) $\frac{1}{(1/30)-0} \left[-\frac{1}{30\pi} \cos(60\pi t) + \frac{1}{120\pi} \sin(120\pi t) \right]_0^{1/30} = 30 \left[\left(\frac{1}{30\pi} \right) - \left(-\frac{1}{30\pi} \right) \right] = 0 \text{ amps}$

109. False

$$\int (2x+1)^2 dx = \frac{1}{2} \int (2x+1)^2 2 dx = \frac{1}{6} (2x+1)^3 + C$$

111. True

$$\int_{-10}^{10} (ax^3 + bx^2 + cx + d) dx = \int_{-10}^{10} (ax^3 + cx) dx + \int_{-10}^{10} (bx^2 + d) dx = 0 + 2 \int_0^{10} (bx^2 + d) dx$$

Odd Even

113. True

$$4 \int \sin x \cos x dx = 2 \int \sin 2x dx = -\cos 2x + C$$

115. Let $u = x + h$, then $du = dx$. When $x = a$, $u = a + h$. When $x = b$, $u = b + h$. Thus,

$$\int_a^b f(x+h) dx = \int_{a+h}^{b+h} f(u) du = \int_{a+h}^{b+h} f(x) dx.$$

Section 4.6 Numerical Integration

1. Exact: $\int_0^2 x^2 dx = \left[\frac{1}{3} x^3 \right]_0^2 = \frac{8}{3} \approx 2.6667$

Trapezoidal: $\int_0^2 x^2 dx \approx \frac{1}{4} \left[0 + 2\left(\frac{1}{2}\right)^2 + 2(1)^2 + 2\left(\frac{3}{2}\right)^2 + (2)^2 \right] = \frac{11}{4} = 2.7500$

Simpson's: $\int_0^2 x^2 dx \approx \frac{1}{6} \left[0 + 4\left(\frac{1}{2}\right)^2 + 2(1)^2 + 4\left(\frac{3}{2}\right)^2 + (2)^2 \right] = \frac{8}{3} \approx 2.6667$

3. Exact: $\int_0^2 x^3 dx = \left[\frac{x^4}{4} \right]_0^2 = 4.000$

Trapezoidal: $\int_0^2 x^3 dx \approx \frac{1}{4} \left[0 + 2\left(\frac{1}{2}\right)^3 + 2(1)^3 + 2\left(\frac{3}{2}\right)^3 + (2)^3 \right] = \frac{17}{4} = 4.2500$

Simpson's: $\int_0^2 x^3 dx \approx \frac{1}{6} \left[0 + 4\left(\frac{1}{2}\right)^3 + 2(1)^3 + 4\left(\frac{3}{2}\right)^3 + (2)^3 \right] = \frac{24}{6} = 4.0000$

5. Exact: $\int_0^2 x^3 dx = \left[\frac{1}{4}x^4 \right]_0^2 = 4.0000$

Trapezoidal: $\int_0^2 x^3 dx \approx \frac{1}{8} \left[0 + 2\left(\frac{1}{4}\right)^3 + 2\left(\frac{2}{4}\right)^3 + 2\left(\frac{3}{4}\right)^3 + 2(1)^3 + 2\left(\frac{5}{4}\right)^3 + 2\left(\frac{6}{4}\right)^3 + 2\left(\frac{7}{4}\right)^3 + 8 \right] = 4.0625$

Simpson's: $\int_0^2 x^3 dx \approx \frac{1}{12} \left[0 + 4\left(\frac{1}{4}\right)^3 + 2\left(\frac{2}{4}\right)^3 + 4\left(\frac{3}{4}\right)^3 + 2(1)^3 + 4\left(\frac{5}{4}\right)^3 + 2\left(\frac{6}{4}\right)^3 + 4\left(\frac{7}{4}\right)^3 + 8 \right] = 4.0000$

7. Exact: $\int_4^9 \sqrt{x} dx = \left[\frac{2}{3}x^{3/2} \right]_4^9 = 18 - \frac{16}{3} = \frac{38}{3} \approx 12.6667$

Trapezoidal: $\int_4^9 \sqrt{x} dx \approx \frac{5}{16} \left[2 + 2\sqrt{\frac{37}{8}} + 2\sqrt{\frac{21}{4}} + 2\sqrt{\frac{47}{8}} + 2\sqrt{\frac{26}{4}} + 2\sqrt{\frac{57}{8}} + 2\sqrt{\frac{31}{4}} + 2\sqrt{\frac{67}{8}} + 3 \right] \approx 12.6640$

Simpson's: $\int_4^9 \sqrt{x} dx \approx \frac{5}{24} \left[2 + 4\sqrt{\frac{37}{8}} + \sqrt{21} + 4\sqrt{\frac{47}{8}} + \sqrt{26} + 4\sqrt{\frac{57}{8}} + \sqrt{31} + 4\sqrt{\frac{67}{8}} + 3 \right] \approx 12.6667$

9. Exact: $\int_1^2 \frac{1}{(x+1)^2} dx = \left[-\frac{1}{x+1} \right]_1^2 = -\frac{1}{3} + \frac{1}{2} = \frac{1}{6} \approx 0.1667$

Trapezoidal: $\int_1^2 \frac{1}{(x+1)^2} dx \approx \frac{1}{8} \left[\frac{1}{4} + 2\left(\frac{1}{((5/4)+1)^2}\right) + 2\left(\frac{1}{((3/2)+1)^2}\right) + 2\left(\frac{1}{((7/4)+1)^2}\right) + \frac{1}{9} \right] = \frac{1}{8} \left(\frac{1}{4} + \frac{32}{81} + \frac{8}{25} + \frac{32}{121} + \frac{1}{9} \right) \approx 0.1676$

Simpson's: $\int_1^2 \frac{1}{(x+1)^2} dx \approx \frac{1}{12} \left[\frac{1}{4} + 4\left(\frac{1}{((5/4)+1)^2}\right) + 2\left(\frac{1}{((3/2)+1)^2}\right) + 4\left(\frac{1}{((7/4)+1)^2}\right) + \frac{1}{9} \right] = \frac{1}{12} \left(\frac{1}{4} + \frac{64}{81} + \frac{8}{25} + \frac{64}{121} + \frac{1}{9} \right) \approx 0.1667$

11. Trapezoidal: $\int_0^2 \sqrt{1+x^3} dx \approx \frac{1}{4} [1 + 2\sqrt{1+(1/8)} + 2\sqrt{2} + 2\sqrt{1+(27/8)} + 3] \approx 3.283$

Simpson's: $\int_0^2 \sqrt{1+x^3} dx \approx \frac{1}{6} [1 + 4\sqrt{1+(1/8)} + 2\sqrt{2} + 4\sqrt{1+(27/8)} + 3] \approx 3.240$

Graphing utility: 3.241

13. $\int_0^1 \sqrt{x} \sqrt{1-x} dx = \int_0^1 \sqrt{x(1-x)} dx$

Trapezoidal: $\int_0^1 \sqrt{x(1-x)} dx \approx \frac{1}{8} \left[0 + 2\sqrt{\frac{1}{4}\left(1-\frac{1}{4}\right)} + 2\sqrt{\frac{1}{2}\left(1-\frac{1}{2}\right)} + 2\sqrt{\frac{3}{4}\left(1-\frac{3}{4}\right)} \right] \approx 0.342$

Simpson's: $\int_0^1 \sqrt{x(1-x)} dx \approx \frac{1}{12} \left[0 + 4\sqrt{\frac{1}{4}\left(1-\frac{1}{4}\right)} + 2\sqrt{\frac{1}{2}\left(1-\frac{1}{2}\right)} + 4\sqrt{\frac{3}{4}\left(1-\frac{3}{4}\right)} \right] \approx 0.372$

Graphing utility: 0.393

15. Trapezoidal: $\int_0^{\sqrt{\pi/2}} \cos(x^2) dx \approx \frac{\sqrt{\pi/2}}{8} \left[\cos 0 + 2 \cos\left(\frac{\sqrt{\pi/2}}{4}\right)^2 + 2 \cos\left(\frac{\sqrt{\pi/2}}{2}\right)^2 + 2 \cos\left(\frac{\sqrt{\pi/2}}{4}\right)^2 + \cos\left(\sqrt{\frac{\pi}{2}}\right)^2 \right] \approx 0.957$

Simpson's: $\int_0^{\sqrt{\pi/2}} \cos(x^2) dx \approx \frac{\sqrt{\pi/2}}{12} \left[\cos 0 + 4 \cos\left(\frac{\sqrt{\pi/2}}{4}\right)^2 + 2 \cos\left(\frac{\sqrt{\pi/2}}{2}\right)^2 + 4 \cos\left(\frac{\sqrt{\pi/2}}{4}\right)^2 + \cos\left(\sqrt{\frac{\pi}{2}}\right)^2 \right] \approx 0.978$

Graphing utility: 0.977

17. Trapezoidal: $\int_1^{1.1} \sin x^2 dx \approx \frac{1}{80} [\sin(1) + 2 \sin(1.025)^2 + 2 \sin(1.05)^2 + 2 \sin(1.075)^2 + \sin(1.1)^2] \approx 0.089$


Simpson's: $\int_1^{1.1} \sin x^2 dx \approx \frac{1}{120} [\sin(1) + 4 \sin(1.025)^2 + 2 \sin(1.05)^2 + 4 \sin(1.075)^2 + \sin(1.1)^2] \approx 0.089$

Graphing utility: 0.089

19. Trapezoidal: $\int_0^{\pi/4} x \tan x dx \approx \frac{\pi}{32} \left[0 + 2\left(\frac{\pi}{16}\right) \tan\left(\frac{\pi}{16}\right) + 2\left(\frac{2\pi}{16}\right) \tan\left(\frac{2\pi}{16}\right) + 2\left(\frac{3\pi}{16}\right) \tan\left(\frac{3\pi}{16}\right) + \frac{\pi}{4} \right] \approx 0.194$

Simpson's: $\int_0^{\pi/4} x \tan x dx \approx \frac{\pi}{48} \left[0 + 4\left(\frac{\pi}{16}\right) \tan\left(\frac{\pi}{16}\right) + 2\left(\frac{2\pi}{16}\right) \tan\left(\frac{2\pi}{16}\right) + 4\left(\frac{3\pi}{16}\right) \tan\left(\frac{3\pi}{16}\right) + \frac{\pi}{4} \right] \approx 0.186$

Graphing utility: 0.186



The Trapezoidal Rule overestimates the area if the graph of the integrand is concave up.

23. $f(x) = x^3$

$f'(x) = 3x^2$

$f''(x) = 6x$

$f'''(x) = 6$

$f^{(4)}(x) = 0$

(a) Trapezoidal: Error $\frac{(2-0)^3}{12(4^2)}(12) = 0.5$ since

$f''(x)$ is maximum in $[0, 2]$ when $x = 2$.

(b) Simpson's: Error $\frac{(2-0)^5}{180(4^4)}(0) = 0$ since

$f^{(4)}(x) = 0$.

25. $f''(x) = \frac{2}{x^3}$ in $[1, 3]$.

(a) $|f''(x)|$ is maximum when $x = 1$ and $|f''(1)| = 2$.

Trapezoidal: Error $\frac{2^3}{12n^2}(2) < 0.00001$, $n^2 > 133,333.33$, $n > 365.15$; let $n = 366$.

$f^{(4)}(x) = \frac{24}{x^5}$ in $[1, 3]$

(b) $|f^{(4)}(x)|$ is maximum when $x = 1$ and when $|f^{(4)}(1)| = 24$.

Simpson's: Error $\frac{2^5}{180n^4}(24) < 0.00001$, $n^4 > 426,666.67$, $n > 25.56$; let $n = 26$.

27. $f(x) = \sqrt{1+x}$

(a) $f''(x) = -\frac{1}{4(1+x)^{3/2}}$ in $[0, 2]$.

$|f''(x)|$ is maximum when $x = 0$ and $|f''(0)| = \frac{1}{4}$.

Trapezoidal: Error $\frac{8}{12n^2} \left(\frac{1}{4}\right) < 0.00001$, $n^2 > 16,666.67$, $n > 129.10$; let $n = 130$.

(b) $f^{(4)}(x) = \frac{-15}{16(1+x)^{7/2}}$ in $[0, 2]$

$|f^{(4)}(x)|$ is maximum when $x = 0$ and $|f^{(4)}(0)| = \frac{15}{16}$.

Simpson's: Error $\frac{32}{180n^4} \left(\frac{15}{16}\right) < 0.00001$, $n^4 > 16,666.67$, $n > 11.36$; let $n = 12$.

29. $f(x) = \tan(x^2)$

(a) $f''(x) = 2 \sec^2(x^2)[1 + 4x^2 \tan(x^2)]$ in $[0, 1]$.

$|f''(x)|$ is maximum when $x = 1$ and $|f''(1)| \approx 49.5305$.

Trapezoidal: Error $\frac{(1-0)^3}{12n^2}(49.5305) < 0.00001$, $n^2 > 412,754.17$, $n > 642.46$; let $n = 643$.

(b) $f^{(4)}(x) = 8 \sec^2(x^2)[12x^2 + (3 + 32x^4) \tan(x^2) + 36x^2 \tan^2(x^2) + 48x^4 \tan^3(x^2)]$ in $[0, 1]$

$|f^{(4)}(x)|$ is maximum when $x = 1$ and $|f^{(4)}(1)| \approx 9184.4734$.

Simpson's: Error $\frac{(1-0)^5}{180n^4}(9184.4734) < 0.00001$, $n^4 > 5,102,485.22$, $n > 47.53$; let $n = 48$.

31. Let $f(x) = Ax^3 + Bx^2 + Cx + D$. Then $f^{(4)}(x) = 0$.

Simpson's: Error $\frac{(b-a)^5}{180n^4}(0) = 0$

Therefore, Simpson's Rule is exact when approximating the integral of a cubic polynomial.

Example: $\int_0^1 x^3 dx = \frac{1}{6} \left[0 + 4 \left(\frac{1}{2}\right)^3 + 1 \right] = \frac{1}{4}$

This is the exact value of the integral.

33. $f(x) = \sqrt{2+3x^2}$ on $[0, 4]$.

n	$L(n)$	$M(n)$	$R(n)$	$T(n)$	$S(n)$
4	12.7771	15.3965	18.4340	15.6055	15.4845
8	14.0868	15.4480	16.9152	15.5010	15.4662
10	14.3569	15.4544	16.6197	15.4883	15.4658
12	14.5386	15.4578	16.4242	15.4814	15.4657
16	14.7674	15.4613	16.1816	15.4745	15.4657
20	14.9056	15.4628	16.0370	15.4713	15.4657


35. $f(x) = \sin \sqrt{x}$ on $[0, 4]$.

n	$L(n)$	$M(n)$	$R(n)$	$T(n)$	$S(n)$
4	2.8163	3.5456	3.7256	3.2709	3.3996
8	3.1809	3.5053	3.6356	3.4083	3.4541
10	3.2478	3.4990	3.6115	3.4296	3.4624
12	3.2909	3.4952	3.5940	3.4425	3.4674
16	3.3431	3.4910	3.5704	3.4568	3.4730
20	3.3734	3.4888	3.5552	3.4643	3.4759

37. $A = \int_0^{\pi/2} \sqrt{x} \cos x \, dx$

Simpson's Rule: $n = 14$

$$\begin{aligned} \int_0^{\pi/2} \sqrt{x} \cos x \, dx &\approx \frac{\pi}{84} \left[\sqrt{0} \cos 0 + 4 \sqrt{\frac{\pi}{28}} \cos \frac{\pi}{28} + 2 \sqrt{\frac{\pi}{14}} \cos \frac{\pi}{14} + 4 \sqrt{\frac{3\pi}{28}} \cos \frac{3\pi}{28} + \dots + \sqrt{\frac{\pi}{2}} \cos \frac{\pi}{2} \right] \\ &\approx 0.701 \end{aligned}$$



39. $W = \int_0^5 100x \sqrt{125 - x^3} \, dx$

Simpson's Rule: $n = 12$

$$\begin{aligned} \int_0^5 100x \sqrt{125 - x^3} \, dx &\approx \frac{5}{3(12)} \left[0 + 400 \left(\frac{5}{12} \right) \sqrt{125 - \left(\frac{5}{12} \right)^3} + 200 \left(\frac{10}{12} \right) \sqrt{125 - \left(\frac{10}{12} \right)^3} \right. \\ &\quad \left. + 400 \left(\frac{15}{12} \right) \sqrt{125 - \left(\frac{15}{12} \right)^3} + \dots + 0 \right] \approx 10,233.58 \text{ ft} \cdot \text{lb} \end{aligned}$$

41. $\int_0^{1/2} \frac{6}{\sqrt{1-x^2}} \, dx$ Simpson's Rule, $n = 6$

$$\begin{aligned} \pi &\approx \frac{\left(\frac{1}{2} - 0 \right)}{3(6)} [6 + 4(6.0209) + 2(6.0851) + 4(6.1968) + 2(6.3640) + 4(6.6002) + 6.9282] \\ &\approx \frac{1}{36} [113.098] \approx 3.1416 \end{aligned}$$

43. Area $\approx \frac{1000}{2(10)} [125 + 2(125) + 2(120) + 2(112) + 2(90) + 2(90) + 2(95) + 2(88) + 2(75) + 2(35)] = 89,250 \text{ sq m}$

45. $\int_0^t \sin \sqrt{x} \, dx = 2, n = 10$

By trial and error, we obtain $t \approx 2.477$.